

LINEAR ALGEBRA

PREVIOUS YEAR PAPERS

DEC – 2014

PART – B

1. Let A, B be $n \times n$ matrices such that $BA + B^2 = I - BA^2$, where I is the $n \times n$ identity matrix. Which of the following is always true?

1. A is nonsingular
2. B is nonsingular
3. A+B is nonsingular
4. AB is nonsingular

2. Which of the following matrices has the

same row space as the matrix $\begin{pmatrix} 4 & 8 & 4 \\ 3 & 6 & 1 \\ 2 & 4 & 0 \end{pmatrix}$?

1. $\begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix}$
2. $\begin{pmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$
3. $\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$
4. $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$

3. The determinant of the $n \times n$ permutation

$$\text{matrix } \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & \ddots & \\ & & & 1 \end{bmatrix}$$

1. $(-1)^n$
2. $(-1)^{\lfloor \frac{n}{2} \rfloor}$
3. -1
4. 1

4. The determinant $\begin{vmatrix} 1 & 1+x & 1+x+x^2 \\ 1 & 1+y & 1+y+y^2 \\ 1 & 1+z & 1+z+z^2 \end{vmatrix}$ is equal to

1. $(z-y)(z-x)(y-x)$
2. $(x-y)(x-z)(y-z)$
3. $(x-y)^2(y-z)^2(z-x)^2$
4. $(x^2-y^2)(y^2-z^2)(z^2-x^2)$

5. Which of the following matrices is not diagonalizable over \mathbb{R} ?

1. $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

2. $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 1 \\ 0 & 0 & 3 \end{bmatrix}$

3. $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}$

4. $\begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$

6. Let P be a 2×2 complex matrix such that P^*P is the identity matrix, where P^* is the conjugate transpose of P. Then the eigenvalues of P are
1. real
 2. complex conjugates of each other
 3. reciprocals of each other
 4. of modulus 1

PART – C

7. Let A be a real $n \times n$ orthogonal matrix, that is, $A^t A = AA^t = I_n$, the $n \times n$ identity matrix. Which of the following statements are necessarily true?

1. $\langle Ax, Ay \rangle = \langle x, y \rangle \forall x, y \in \mathbb{R}^n$
2. All eigenvalues of A are either +1 or -1.
3. The rows of A form an orthonormal basis of \mathbb{R}^n
4. A is diagonalizable over \mathbb{R} .

8. Which of the following matrices have Jordan canonical form equal to $\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$

1. $\begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$

2. $\begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$

3. $\begin{pmatrix} 0 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$

4. $\begin{pmatrix} 0 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$

9. Let A be a 3×4 and b be a 3×1 matrix with integer entries. Suppose that the system $Ax=b$ has a complex solution. Then
1. $Ax=b$ has an integer solution
 2. $Ax=b$ has a rational solution
 3. The set of real solutions to $Ax=0$ has a basis consisting of rational solutions.
 4. If $b \neq 0$, then A has positive rank.

10. Let f be a non-zero symmetric bilinear form on \mathbb{R}^3 . Suppose that there exists linear transformations $T_i: \mathbb{R}^3 \rightarrow \mathbb{R}$, $i = 1, 2$ such that for all $\alpha, \beta \in \mathbb{R}^3$, $f(\alpha, \beta) = T_1(\alpha) T_2(\beta)$. Then
- rank $f=1$
 - $\dim \{\beta \in \mathbb{R}^3 : f(\alpha, \beta) = 0 \text{ for all } \alpha \in \mathbb{R}^3\} = 2$
 - f is positive semi-definite or negative semi-definite.
 - $\{\alpha : f(\alpha, \alpha) = 0\}$ is a linear subspace of dimension 2

11. The matrix $A = \begin{pmatrix} 5 & 9 & 8 \\ 1 & 8 & 2 \\ 9 & 1 & 0 \end{pmatrix}$ satisfies

- A is invertible and the inverse has all integer entries.
- $\det(A)$ is odd.
- $\det(A)$ is divisible by 13.
- $\det(A)$ has at least two prime divisors.

12. Let A be 5×5 matrix and let B be obtained by changing one element of A . Let r and s be the ranks of A and B respectively. Which of the following statements is/are correct?
- $s \leq r+1$
 - $r-1 \leq s$
 - $s = r-1$
 - $s \neq r$

13. Let $M_n(K)$ denote the space of all $n \times n$ matrices with entries in a field K . Fix a non-singular matrix $A = (A_{ij}) \in M_n(K)$, and consider the linear map $T: M_n(K) \rightarrow M_n(K)$ given by $T(X) = AX$. Then
- $\text{trace}(T) = n \sum_{i=1}^n A_{ii}$
 - $\text{trace}(T) = \sum_{i=1}^n \sum_{j=1}^n A_{ij}$
 - rank of T is n^2
 - T is non-singular

14. For arbitrary subspaces U, V and W of a finite dimensional vector space, which of the following hold
- $U \cap (V+W) \subset U \cap V + U \cap W$
 - $U \cap (V+W) \supset U \cap V + U \cap W$
 - $(U \cap V) + W \subset (U+W) \cap (V+W)$
 - $(U \cap V) + W \supset (U+W) \cap (V+W)$

15. Let A be 4×7 real matrix and B be a 7×4 real matrix such that $AB = I_4$, where I_4 is

the 4×4 identity matrix. Which of the following is/are always true?

- rank $(A)=4$
- rank $(B)=7$
- nullity $(B)=0$
- $BA = I_7$, where I_7 is the 7×7 identity matrix

16. Let $\mathbb{R}[x]$ denote the vector space of all real polynomials. Let $D: \mathbb{R}[x] \rightarrow \mathbb{R}[x]$ denote the map $Df = \frac{df}{dx}, \forall f$. Then,
- D is one-one
 - D is onto
 - There exists $E: \mathbb{R}[x] \rightarrow \mathbb{R}[x]$ so that $D(E(f)) = f, \forall f$.
 - There exists $E: \mathbb{R}[x] \rightarrow \mathbb{R}[x]$ so that $E(D(f)) = f, \forall f$.

17. Which of the following are eigenvalues of the

matrix $\begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix} ?$

- +1
- 1
- +i
- i

18. Let $A = \begin{pmatrix} x & y \\ -y & x \end{pmatrix}$, where $x, y \in \mathbb{R}$ such that $x^2 + y^2 = 1$. Then we must have

- For any $n \geq 1$, $A^n = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$ where $x = \cos(\theta/n), y = \sin(\theta/n)$
- $\text{tr}(A) \neq 0$
- $A^t = A^{-1}$
- A is similar to a diagonal matrix over \mathbb{C}

JUNE – 2015

PART – B

19. Let V be the space of twice differentiable functions on \mathbb{R} satisfying $f'' - 2f' + f = 0$.

Define $T: V \rightarrow \mathbb{R}^2$ by $T(f) = (f'(0), f(0))$.

Then T is

1. one-to-one and onto
2. one-to-one but not onto
3. onto but not one-to-one
4. neither one-to-one nor onto

20. The row space of a 20×50 matrix A has dimension 13. What is the dimension of the space of solutions of $Ax = 0$?

1. 7
2. 13
3. 33
4. 37

21. Let A, B be $n \times n$ matrices. Which of the following equals $\text{trace}(A^2B^2)$?

1. $(\text{trace}(AB))^2$
2. $\text{trace}(AB^2A)$
3. $\text{trace}((AB)^2)$
4. $\text{trace}(BABA)$

22. Given a 4×4 real matrix A, let $T: \mathbb{R}^4 \rightarrow \mathbb{R}^4$ be the linear transformation defined by $Tv = Av$, where we think of \mathbb{R}^4 as the set of real 4×1 matrices. For which choices of A given below, do $\text{Image}(T)$ and $\text{Image}(T^2)$ have respective dimensions 2 and 1? (* denotes a non zero entry)

1. $A = \begin{bmatrix} 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & 0 \end{bmatrix}$

2. $A = \begin{bmatrix} 0 & 0 & * & 0 \\ 0 & 0 & * & 0 \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \end{bmatrix}$

3. $A = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & * \\ 0 & 0 & * & 0 \end{bmatrix}$

4. $A = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & * & * \\ 0 & 0 & * & * \end{bmatrix}$

23. Let T be a 4×4 real matrix such that $T^4 = 0$. Let $k_i = \dim \text{Ker} T^i$ for $1 \leq i \leq 4$. Which of the following is NOT a possibility for the sequence $k_1 \leq k_2 \leq k_3 \leq k_4$?

1. $3 \leq 4 \leq 4 \leq 4$
2. $1 \leq 3 \leq 4 \leq 4$
3. $2 \leq 4 \leq 4 \leq 4$
4. $2 \leq 3 \leq 4 \leq 4$

24. Which of the following is a linear transformation from \mathbb{R}^3 to \mathbb{R}^2 ?

(a) $f \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 4 \\ x+y \end{pmatrix}$ (b) $g \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} xy \\ x+y \end{pmatrix}$

(c) $h \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} z-x \\ x+y \end{pmatrix}$

1. Only f.
2. Only g.
3. Only h.
4. all the transformations f, g and h.

PART - C

25. Let A be an $m \times n$ matrix of rank n with real entries. Choose the correct statement.

1. $Ax = b$ has a solution for any b.
2. $Ax = 0$ does not have a solution.
3. If $Ax = b$ has a solution, then it is unique.
4. $y'A = 0$ for some nonzero y, where y' denotes the transpose of the vector y.

26. Let $F: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ be the function $F(x, y) = \langle Ax, y \rangle$ where $\langle \cdot, \cdot \rangle$ is the standard

inner product of \mathbb{R}^n and A is a $n \times n$ real matrix. Here D denotes the total derivative. Which of the following statements are correct?

1. $(DF(x, y))(u, v) = \langle Au, y \rangle + \langle Ax, v \rangle$
2. $(DF(x, y))(0, 0) = 0$.
3. $DF(x, y)$ may not exist for some $(x, y) \in \mathbb{R}^n \times \mathbb{R}^n$
4. $DF(x, y)$ does not exist at $(x, y) = (0, 0)$.

27. Let $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a continuous function such that $\int_{\mathbb{R}^n} |f(x)| dx < \infty$. Let A be a real $n \times n$

invertible matrix and for $x, y \in \mathbb{R}^n$, let $\langle x, y \rangle$ denote the standard inner product in \mathbb{R}^n .

Then $\int_{\mathbb{R}^n} f(Ax) e^{i\langle y, x \rangle} dx =$

1. $\int_{\mathbb{R}^n} f(x) e^{i\langle (A^{-1})^T y, x \rangle} \frac{dx}{|\det A|}$
2. $\int_{\mathbb{R}^n} f(x) e^{i\langle A^T y, x \rangle} \frac{dx}{|\det A|}$
3. $\int_{\mathbb{R}^n} f(x) e^{i\langle (A^T)^{-1} y, x \rangle} dx$

$$4. \int_{\mathbb{R}^n} f(x) e^{i(A^{-1}y \cdot x)} \frac{dx}{|\det A|}$$

28. Let S be the set of 3x3 real matrices A with

$$A^T A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}. \text{ Then the set S contains}$$

1. a nilpotent matrix.
 2. a matrix of rank one.
 3. a matrix of rank two.
 4. a non-zero skew-symmetric matrix.
29. An nxn complex matrix A satisfies $A^k = I_n$, the nxn identity matrix, where k is a positive integer > 1 . Suppose 1 is not an eigenvalue of A. Then which of the following statements are necessarily true?

1. A is diagonalizable.
2. $A + A^2 + \dots + A^{k-1} = O$, the nxn zero matrix
3. $\text{tr}(A) + \text{tr}(A^2) + \dots + \text{tr}(A^{k-1}) = -n$
4. $A^{-1} + A^{-2} + \dots + A^{-(k-1)} = -I_n$

30. Let $S: \mathbb{R}^n \rightarrow \mathbb{R}^n$ be given by $S(v) = \alpha v$ for a fixed $\alpha \in \mathbb{R}, \alpha \neq 0$.

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a linear transformation such that $B = \{v_1, \dots, v_n\}$ is a set of linearly independent eigen vectors of T. Then

1. The matrix of T with respect to B is diagonal.
2. The matrix of (T - S) with respect to B is diagonal.
3. The matrix of T with respect to B is not necessarily diagonal, but is upper triangular.
4. The matrix of T with respect to B is diagonal but the matrix of (T-S) with respect to B is not diagonal.

31. Let $p_n(x) = x^n$ for $x \in \mathbb{R}$ and let $\wp = \text{span}\{p_0, p_1, p_2, \dots\}$. Then

1. \wp is the vector space of all real valued continuous functions on \mathbb{R} .
2. \wp is a subspace of all real valued continuous functions on \mathbb{R} .
3. $\{p_0, p_1, p_2, \dots\}$ is a linearly independent set in vector space of all continuous functions on \mathbb{R} .
4. Trigonometric functions belong to \wp .

32. Let $A = \begin{bmatrix} a & b & c \\ 0 & a & d \\ 0 & 0 & a \end{bmatrix}$ be a 3x3 matrix where

a,b,c,d are integers. Then, we must have:

1. If $a \neq 0$, there is a polynomial $p \in \mathbb{Q}[x]$ such that $p(A)$ is the inverse of A.

2. For each polynomial $q \in \mathbb{Z}[x]$, the matrix

$$q(A) = \begin{bmatrix} q(a) & q(b) & q(c) \\ 0 & q(a) & q(d) \\ 0 & 0 & q(a) \end{bmatrix}$$

3. If $A^n = O$ for some positive integer n, then $A^3 = O$.

4. A commutes with every matrix of the

$$\text{form } \begin{bmatrix} a' & 0 & c' \\ 0 & a' & 0 \\ 0 & 0 & a' \end{bmatrix}.$$

33. Which of the following are subspaces of vector space \mathbb{R}^3 ?

1. $\{(x,y,z) : x + y = 0\}$
2. $\{(x,y,z) : x - y = 0\}$
3. $\{(x,y,z) : x + y = 1\}$
4. $\{(x,y,z) : x - y = 1\}$

34. Consider non-zero vector spaces V_1, V_2, V_3, V_4 and linear transformations $T_1: V_1 \rightarrow V_2, T_2: V_2 \rightarrow V_3, T_3: V_3 \rightarrow V_4$ such that $\text{Ker}(T_1) = \{0\}, \text{Range}(T_1) = \text{Ker}(T_2), \text{Range}(T_2) = \text{Ker}(T_3), \text{Range}(T_3) = V_4$. Then

1. $\sum_{i=1}^4 (-1)^i \dim V_i = 0$
2. $\sum_{i=2}^4 (-1)^i \dim V_i > 0$
3. $\sum_{i=1}^4 (-1)^i \dim V_i < 0$
4. $\sum_{i=1}^4 (-1)^i \dim V_i \neq 0$

35. Let A be an invertible 4x4 real matrix. Which of the following are NOT true?

1. Rank A = 4.
2. For every vector $b \in \mathbb{R}^4, Ax = b$ has exactly one solution.
3. $\dim(\text{nullspace } A) \geq 1$.
4. 0 is an eigenvalue of A.

36. Let \underline{u} be a real nx1 vector satisfying $\underline{u}'\underline{u}=1$, where \underline{u}' is the transpose of \underline{u} . Define $A = I - 2\underline{u}\underline{u}'$ where I is the nth order identity matrix. Which of the following statements are true?

1. A is singular
2. $A^2 = A$
3. $\text{Trace}(A) = n-2$
4. $A^2 = I$

DEC – 2015

PART – B

37. Let S denote the set of all the prime numbers p with the property that the matrix

$$\begin{bmatrix} 91 & 31 & 0 \\ 29 & 31 & 0 \\ 79 & 23 & 59 \end{bmatrix} \text{ has an inverse in the field}$$

$\mathbb{Z}/p\mathbb{Z}$. Then

1. $S = \{31\}$
 2. $S = \{31, 59\}$
 3. $S = \{7, 13, 59\}$
 4. S is infinite
38. For a positive integer n , let P_n denote the vector space of polynomials in one variable x with real coefficients and with degree $\leq n$. Consider the map $T: P_2 \rightarrow P_4$ defined by $T(p(x)) = p(x^2)$. Then
1. T is a linear transformation and $\dim \text{range}(T) = 5$.
 2. T is a linear transformation and $\dim \text{range}(T) = 3$.
 3. T is a linear transformation and $\dim \text{range}(T) = 2$.
 4. T is not a linear transformation.
39. Let A be a real 3×4 matrix of rank 2. Then the rank of $A^t A$, where A^t denotes the transpose of A , is:
1. exactly 2
 2. exactly 3
 3. exactly 4
 4. at most 2 but not necessarily 2
40. Consider the quadratic form $Q(v) = v^t A v$,
- $$\text{where } A = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}, v = (x, y, z, w)$$
- Then
1. Q has rank 3.
 2. $xy + z^2 = Q(Pv)$ for some invertible 4×4 real matrix P
 3. $xy + y^2 + z^2 = Q(Pv)$ for some invertible 4×4 real matrix P
 4. $x^2 + y^2 - zw = Q(Pv)$ for some invertible 4×4 real matrix P .

41. If A is a 5×5 real matrix with trace 15 and if 2 and 3 are eigenvalues of A , each with algebraic multiplicity 2, then the determinant of A is equal to
1. 0
 2. 24
 3. 120
 4. 180

42. Let $A \neq I_n$ be an $n \times n$ matrix such that $A^2 = A$, where I_n is the identity matrix of order n . Which of the following statements is false?
1. $(I_n - A)^2 = I_n - A$.
 2. $\text{Trace}(A) = \text{Rank}(A)$.
 3. $\text{Rank}(A) + \text{Rank}(I_n - A) = n$.
 4. The eigenvalues of A are each equal to 1.

PART – C

43. Let A and B be $n \times n$ matrices over \mathbb{C} . Then,
1. AB and BA always have the same set of eigenvalues.
 2. If AB and BA have the same set of eigenvalues then $AB = BA$.
 3. If A^{-1} exists then AB and BA are similar.
 4. The rank of AB is always the same as the rank of BA .
44. Let A be an $m \times n$ real matrix and $b \in \mathbb{R}^m$ with $b \neq 0$.
1. The set of all real solutions of $Ax = b$ is a vector space.
 2. If u and v are two solutions of $Ax = b$, then $\lambda u + (1 - \lambda)v$ is also a solution of $Ax = b$, for any $\lambda \in \mathbb{R}$.
 3. For any two solutions u and v of $Ax = b$, the linear combination $\lambda u + (1 - \lambda)v$ is also a solution of $Ax = b$ only when $0 \leq \lambda \leq 1$.
 4. If rank of A is n , then $Ax = b$ has at most one solution.
45. Let A be an $n \times n$ matrix over \mathbb{C} such that every nonzero vector of \mathbb{C}^n is an eigenvector of A . Then.
1. All eigenvalues of A are equal.
 2. All eigenvalues of A are distinct.
 3. $A = \lambda I$ for some $\lambda \in \mathbb{C}$, where I is the $n \times n$ identity matrix.
 4. If χ_A and m_A denote the characteristic polynomial and the minimal polynomial respectively, then $\chi_A = m_A$.

46. Consider the matrices $A = \begin{bmatrix} 2 & 2 & 1 \\ 0 & 2 & -1 \\ 0 & 0 & 3 \end{bmatrix}$ and

$$B = \begin{bmatrix} 2 & 1 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}.$$

Then

1. A and B are similar over the field of rational numbers \mathbb{Q} .

2. A is diagonalizable over the field of rational numbers \mathbb{Q} .
3. B is the Jordan canonical form of A.
4. The minimal polynomial and the characteristic polynomial of A are the same
47. Let V be a finite dimensional vector space over \mathbb{R} . Let $T : V \rightarrow V$ be a linear transformation such that $\text{rank}(T^2) = \text{rank}(T)$. Then,
1. $\text{Kernel}(T^2) = \text{Kernel}(T)$.
 2. $\text{Range}(T^2) = \text{Range}(T)$.
 3. $\text{Kernel}(T) \cap \text{Range}(T) = \{0\}$.
 4. $\text{Kernel}(T^2) \cap \text{Range}(T^2) = \{0\}$.
48. Let V be the vector space of polynomials over \mathbb{R} of degree less than or equal to n. For $p(x) = a_0 + a_1x + \dots + a_nx^n$ in V, define a linear transformation $T:V \rightarrow V$ by $(Tp)(x) = a_n + a_{n-1}x + \dots + a_0x^n$. Then
1. T is one to one.
 2. T is onto.
 3. T is invertible.
 4. $\det T = \pm 1$.

JUNE – 2016

PART – B

49. Given a $n \times n$ matrix B define e^B by $e^B = \sum_{j=0}^{\infty} \frac{B^j}{j!}$. Let p be the characteristic polynomial of B. Then the matrix $e^{p(B)}$ is
1. $I_{n \times n}$
 2. $0_{n \times n}$
 3. $eI_{n \times n}$
 4. $\pi I_{n \times n}$
50. Let A be a $n \times m$ matrix and b be a $n \times 1$ vector (with real entries). Suppose the equation $Ax=b$, $x \in \mathbb{R}^m$ admits a unique solution. Then we can conclude that
1. $m \geq n$
 2. $n \geq m$
 3. $n = m$
 4. $n > m$
51. Let V be the vector space of all real polynomials of degree ≤ 10 . Let $Tp(x) = p'(x)$ for $p \in V$ be a linear transformation from V to V. Consider the basis $\{1, x, x^2, \dots, x^{10}\}$ of V. Let A be the matrix of T with respect to this basis. Then
1. $\text{Trace} A = 1$
 2. $\det A = 0$
 3. there is no $m \in \mathbb{N}$ such that $A^m = 0$
 4. A has a non zero eigenvalue

52. Let $x = (x_1, x_2, x_3), y = (y_1, y_2, y_3) \in \mathbb{R}^3$ be linearly independent. Let $\delta_1 = x_2y_3 - y_2x_3$, $\delta_2 = x_1y_3 - y_1x_3$, $\delta_3 = x_1y_2 - y_1x_2$. If V is the span of x,y then
1. $V = \{(u, v, w) : \delta_1u - \delta_2v + \delta_3w = 0\}$
 2. $V = \{(u, v, w) : -\delta_1u + \delta_2v + \delta_3w = 0\}$
 3. $V = \{(u, v, w) : \delta_1u + \delta_2v - \delta_3w = 0\}$
 4. $V = \{(u, v, w) : \delta_1u + \delta_2v + \delta_3w = 0\}$
53. Let A be a $n \times n$ real symmetric non-singular matrix. Suppose there exists $x \in \mathbb{R}^n$ such that $x'Ax < 0$. Then we can conclude that
1. $\det(A) < 0$
 2. B = -A is positive definite
 3. $\exists y \in \mathbb{R}^n; y'A^{-1}y < 0$
 4. $\forall y \in \mathbb{R}^n; y'A^{-1}y < 0$

54. Let $A = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$. Let $f : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by $f(v, w) = w^T Av$. Pick the correct statement from below:
1. There exists an eigenvector v of A such that Av is perpendicular to v
 2. The set $\{v \in \mathbb{R}^2 \mid f(v, v) = 0\}$ is a nonzero subspace of \mathbb{R}^2
 3. If $v, w \in \mathbb{R}^2$ are non zero vectors such that $f(v, v) = 0 = f(w, w)$, then v is a scalar multiple of w.
 4. For every $v \in \mathbb{R}^2$, there exists a non zero $w \in \mathbb{R}^2$ such that $f(v, w) = 0$.

PART – C

55. Let V be the vector space of all complex polynomials p with $\deg p \leq n$. Let $T : V \rightarrow V$ be the map $(Tp)(x) = p'(1)$, $x \in \mathbb{C}$. Which of the following are correct?
1. $\dim \text{Ker } T = n$.
 2. $\dim \text{range } T = 1$.
 3. $\dim \text{Ker } T = 1$.
 4. $\dim \text{range } T = n+1$.
56. Let A be an $n \times n$ real matrix. Pick the correct answer(s) from the following
1. A has at least one real eigenvalue.
 2. For all nonzero vectors $v, w \in \mathbb{R}^n$, $(Aw)^T(Av) > 0$.
 3. Every eigenvalue of $A^T A$ is a non negative real number.
 4. $I + A^T A$ is invertible.

57. Let T be a n x n matrix with the property T^n=0. Which of the following is/are true?

- 1. T has n distinct eigenvalues
2. T has one eigenvalue of multiplicity n
3. 0 is an eigenvalue of T.
4. T is similar to a diagonal matrix.

58. Let V = {f: [0,1] -> R | f is a polynomial of degree less than or equal to n}. Let f_j(x) = x^j for 0 <= j <= n and let A be the (n+1) x (n+1) matrix given by a_ij = integral from 0 to 1 of f_i(x)f_j(x)dx.

Then which of the following is/are true?

- 1. dim V = n.
2. dim V > n.
3. A is nonnegative definite, i.e., for all v in R^n, <Av, v> >= 0.
4. det A > 0.

59. Consider the real vector space V of polynomials of degree less than or equal to d. For p in V define ||p||_k = max { |p(0)|, |p'(0)|, ..., |p^(k)(0)| }, where p^(i)(0) is the i-th derivative of p evaluated at 0. Then ||p||_k defines a norm on V if and only if

- 1. k >= d - 1
2. k < d
3. k >= d
4. k < d - 1

60. Let A, B be nxn real matrices such that det A > 0 and det B < 0. For 0 <= t <= 1. Consider C(t) = tA + (1-t)B. Then

- 1. C(t) is invertible for each t in [0,1].
2. There is a t_0 in (0,1) such that C(t_0) is not invertible.
3. C(t) is not invertible for each t in [0,1].
4. C(t) is invertible for only finitely many t in [0,1].

61. Let {a_1, ..., a_n} and {b_1, ..., b_n} be two bases of R^n. Let P be nxn matrix with real entries such that Pa_i = b_i i=1,2,...,n. Suppose that every eigenvalue of P is either -1 or 1. Let Q = I + 2P.

- Then which of the following statements are true?
1. {a_i + 2b_i | i=1,2,...,n} is also a basis of V.
2. Q is invertible.
3. Every eigenvalue of Q is either 3 or -1.
4. det Q > 0 if det P > 0.

62. Let A be an n x n matrix with real entries. Define <x, y>_A = <Ax, Ay>, x, y in R^n.

Then <x, y>_A defines an inner-product if and only if

- 1. ker A = {0}.
2. rank A = n.
3. All eigenvalues of A are positive.
4. All eigenvalues of A are non-negative.

63. Suppose {v_1, ..., v_n} are unit vectors in R^n

such that ||v||^2 = sum from i=1 to n of |<v_i, v>|^2 for all v in R^n

Then decide the correct statements in the following

- 1. v_1, ..., v_n are mutually orthogonal
2. {v_1, ..., v_n} is a basis for R^n
3. v_1, ..., v_n are not mutually orthogonal
4. At most n - 1 of the elements in the set {v_1, ..., v_n} can be orthogonal.

DEC - 2016

PART - B

64. The matrix [[3 -1 0], [-1 2 -1], [0 -1 3]] is

- 1. positive definite.
2. non-negative definite but not positive definite.
3. negative definite
4. neither negative definite nor positive definite

65. Which of the following subsets of R^4 is a basis of R^4?

- B_1 = {(1,0,0,0), (1,1,0,0), (1,1,1,0), (1,1,1,1)}
B_2 = {(1,0,0,0), (1,2,0,0), (1,2,3,0), (1,2,3,4)}
B_3 = {(1,2,0,0), (0,0,1,1), (2,1,0,0), (-5,5,0,0)}

- 1. B_1 and B_2 but not B_3
2. B_1, B_2 and B_3
3. B_1 and B_3 but not B_2
4. Only B_1

66. Let D_1 = det [[a b c], [x y z], [p q r]] and

D_2 = det [[-x a -p], [y -b q], [z -c r]]. Then

- 1. D_1 = D_2
2. D_1 = 2D_2
3. D_1 = -D_2
4. 2D_1 = D_2

67. Consider the matrix $A = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$,

where $\theta = \frac{2\pi}{31}$. Then A^{2015} equals

1. A
2. I
3. $\begin{pmatrix} \cos 13\theta & \sin 13\theta \\ -\sin 13\theta & \cos 13\theta \end{pmatrix}$
4. $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$

68. Let J denote the matrix of order $n \times n$ with all entries 1 and let B be a $(3n) \times (3n)$ matrix

$$\text{given by } B = \begin{pmatrix} 0 & 0 & J \\ 0 & J & 0 \\ J & 0 & 0 \end{pmatrix}$$

Then the rank of B is

1. $2n$
2. $3n - 1$
3. 2
4. 3

69. Which of the following sets of functions from \mathbb{R} to \mathbb{R} is a vector space over \mathbb{R} ?

$$S_1 = \{f \mid \lim_{x \rightarrow 3} f(x) = 0\}$$

$$S_2 = \left\{g \mid \lim_{x \rightarrow 3} g(x) = 1\right\}$$

$$S_3 = \left\{h \mid \lim_{x \rightarrow 3} h(x) \text{ exists}\right\}$$

1. Only S_1
2. Only S_2
3. S_1 and S_3 but not S_2
4. All the three are vector spaces

70. Let A be an $n \times m$ matrix with each entry equal to +1, -1 or 0 such that every column has exactly one +1 and exactly one -1. We can conclude that

1. Rank $A \leq n - 1$
2. Rank $A = m$
3. $n \leq m$
4. $n - 1 \leq m$

71. What is the number of non-singular 3×3 matrices over F_2 , the finite field with two elements?

1. 168
2. 384
3. 2^3
4. 3^2

PART - C

72. Let $A = [a_{ij}]$ be an $n \times n$ matrix such that a_{ij} is an integer for all i, j. Let $AB = I$ with $B = [b_{ij}]$ (where I is the identity matrix). For a square

matrix C, det C denotes its determinant. Which of the following statements is true?

1. If $\det A = 1$ then $\det B = 1$.
2. A sufficient condition for each b_{ij} to be an integer is that $\det A$ is an integer.
3. B is always an integer matrix.
4. A necessary condition for each b_{ij} to be an integer is $\det A \in \{-1, +1\}$.

73. Let $A = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$ and let α_n and β_n denote

the two eigenvalues of A^n such that $|\alpha_n| \geq |\beta_n|$. Then

1. $\alpha_n \rightarrow \infty$ as $n \rightarrow \infty$
2. $\beta_n \rightarrow 0$ as $n \rightarrow \infty$
3. β_n is positive if n is even.
4. β_n is negative if n is odd.

74. Let M_n denote the vector space of all $n \times n$ real matrices. Among the following subsets of M_n , decide which are linear subspaces.

1. $V_1 = \{A \in M_n : A \text{ is nonsingular}\}$
2. $V_2 = \{A \in M_n : \det(A) = 0\}$
3. $V_3 = \{A \in M_n : \text{trace}(A) = 0\}$
4. $V_4 = \{BA : A \in M_n\}$, where B is some fixed matrix in M_n .

75. If P and Q are invertible matrices such that $PQ = -QP$, then we can conclude that

1. $\text{Tr}(P) = \text{Tr}(Q) = 0$
2. $\text{Tr}(P) = \text{Tr}(Q) = 1$
3. $\text{Tr}(P) = -\text{Tr}(Q)$
4. $\text{Tr}(P) \neq \text{Tr}(Q)$

76. Let n be an odd number ≥ 7 . Let $A = [a_{ij}]$ be an $n \times n$ matrix with $a_{i,i+1} = 1$ for all $i = 1, 2, \dots, n-1$ and $a_{n,1} = 1$. Let $a_{ij} = 0$ for all the other pairs (i, j). Then we can conclude that

1. A has 1 as an eigenvalue.
2. A has -1 as an eigenvalue.
3. A has at least one eigenvalue with multiplicity ≥ 2 .
4. A has no real eigenvalues.

77. Let W_1, W_2, W_3 be three distinct subspaces of \mathbb{R}^{10} such that each W_i has dimension 9. Let $W = W_1 \cap W_2 \cap W_3$. Then we can conclude that

1. W may not be a subspace of \mathbb{R}^{10}
2. $\dim W \leq 8$
3. $\dim W \geq 7$
4. $\dim W \leq 3$

78. Let A be a real symmetric matrix. Then we can conclude that
1. A does not have 0 as an eigenvalue
 2. All eigenvalues of A are real
 3. If A^{-1} exists, then A^{-1} is real and symmetric
 4. A has at least one positive eigenvalue

JUNE- 2017

PART - B

79. Let A be a 4×4 matrix. Suppose that the null space $N(A)$ of A is $\{(x, y, z, w) \in \mathbb{R}^4 : x+y+z = 0, x+y+w = 0\}$. Then
1. $\dim(\text{column space}(A)) = 1$
 2. $\dim(\text{column space}(A)) = 2$
 3. $\text{rank}(A) = 1$
 4. $S = \{(1,1,1,0), (1,1,0,1)\}$ is a basis of $N(A)$

80. Let A and B be real invertible matrices such that $AB = -BA$. Then
1. $\text{Trace}(A) = \text{Trace}(B) = 0$
 2. $\text{Trace}(A) = \text{Trace}(B) = 1$
 3. $\text{Trace}(A) = 0, \text{Trace}(B) = 1$
 4. $\text{Trace}(A) = 1, \text{Trace}(B) = 0$

81. Let A be an $n \times n$ self-adjoint matrix with eigenvalues $\lambda_1, \dots, \lambda_n$.
Let $\|X\|_2 = \sqrt{|x_1|^2 + \dots + |x_n|^2}$ for $X = (x_1, \dots, x_n) \in \mathbb{C}^n$.
If $p(A) = a_0I + a_1A + \dots + a_nA^n$ then $\sup_{\|X\|_2=1} \|p(A)X\|_2$ is equal to
1. $\max\{a_0 + a_1\lambda_j + \dots + a_n\lambda_j^n : 1 \leq j \leq n\}$
 2. $\max\{|a_0 + a_1\lambda_j + \dots + a_n\lambda_j^n| : 1 \leq j \leq n\}$
 3. $\min\{a_0 + a_1\lambda_j + \dots + a_n\lambda_j^n : 1 \leq j \leq n\}$
 4. $\min\{|a_0 + a_1\lambda_j + \dots + a_n\lambda_j^n| : 1 \leq j \leq n\}$

82. Let $p(x) = \alpha x^2 + \beta x + \gamma$ be a polynomial, where $\alpha, \beta, \gamma \in \mathbb{R}$. Fix $x_0 \in \mathbb{R}$. Let $S = \{(a, b, c) \in \mathbb{R}^3 : p(x) = a(x - x_0)^2 + b(x - x_0) + c \text{ for all } x \in \mathbb{R}\}$.
Then the number of elements in S is
1. 0
 2. 1
 3. strictly greater than 1 but finite
 4. infinite

83. Let $A = \begin{bmatrix} 1 & 0 & 2 \\ 1 & -2 & 0 \\ 0 & 0 & -3 \end{bmatrix}$ and I be the 3×3 identity matrix. If $6A^{-1} = aA^2 + bA + cI$ for $a, b, c \in \mathbb{R}$ then (a, b, c) equals
1. (1, 2, 1)
 2. (1, -1, 2)
 3. (4, 1, 1)
 4. (1, 4, 1)

84. Let $A = \begin{bmatrix} 1 & 1 & 2 \\ 1 & -2 & 5 \\ 2 & 5 & -3 \end{bmatrix}$.
Then the eigenvalues of A are
1. -4, 3, -3
 2. 4, 3, 1
 3. $4, -4 \pm \sqrt{13}$
 4. $4, -2 \pm 2\sqrt{7}$

PART - C

85. Consider the vector space V of real polynomials of degree less than or equal to n . Fix distinct real numbers a_0, a_1, \dots, a_k . For $p \in V$, $\max\{|p(a_j)| : 0 \leq j \leq k\}$ defines a norm on V
1. only if $k < n$
 2. only if $k \geq n$
 3. if $k+1 \leq n$
 4. if $k \geq n+1$
86. Let V be the vector space of polynomials of degree at most 3 in a variable x with coefficient in \mathbb{R} . Let $T = d/dx$ be the linear transformation of V to itself given by differentiation. Which of the following are correct?
1. T is invertible
 2. 0 is an eigenvalue of T
 3. There is a basis with respect to which the matrix of T is nilpotent.
 4. The matrix of T with respect to the basis $\{1, 1+x, 1+x+x^2, 1+x+x^2+x^3\}$ is diagonal
87. Let m, n, r be natural numbers. Let A be $m \times n$ matrix with real entries such that $(AA^t)^r = I$, where I is the $m \times m$ identity matrix and A^t is the transpose of the matrix A . We can conclude that
1. $m=n$
 2. AA^t is invertible
 3. $A^t A$ is invertible
 4. if $m=n$, then A is invertible
88. Let A be an $n \times n$ real matrix with $A^2 = A$. Then
1. the eigenvalues of A are either 0 or 1

- 2. A is a diagonal matrix with diagonal entries 0 or 1
- 3. rank(A) = trace (A)
- 4. rank (I - A) = trace (I - A)

89. For any $n \times n$ matrix B, let $N(B) = \{X \in \mathbb{R}^n : BX = 0\}$ be the null space of B. Let A be a 4×4 matrix with $\dim(N(A - 2I))=2$, $\dim(N(A - 4I))=1$ and rank (A) = 3. Then
- 1. 0, 2 and 4 are eigenvalues of A
 - 2. determinant (A) = 0
 - 3. A is not diagonalizable
 - 4. trace (A) = 8

90. Which of the following 3×3 matrices are diagonalizable over \mathbb{R} ?

- 1. $\begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{bmatrix}$
- 2. $\begin{bmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$
- 3. $\begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 4 \\ 3 & 4 & 1 \end{bmatrix}$
- 4. $\begin{bmatrix} 0 & 1 & 2 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$

91. Let H be a real Hilbert space and $M \subseteq H$ be a closed linear subspace. Let $x_0 \in H \setminus M$. Let $y_0 \in M$ be such that $\|x_0 - y_0\| = \inf\{\|x_0 - y\| : y \in M\}$. Then
- 1. such a y_0 is unique
 - 2. $x_0 \perp M$
 - 3. $y_0 \perp M$
 - 4. $x_0 - y_0 \perp M$

92. Let $A = \begin{bmatrix} 3 & 1 & 2 \\ 1 & 2 & 3 \\ 2 & 3 & 1 \end{bmatrix}$ and $Q(X) = X^T A X$ for

- $X \in \mathbb{R}^3$. Then
- 1. A has exactly two positive eigenvalues
 - 2. all the eigenvalues of A are positive
 - 3. $Q(X) \geq 0$ for all $X \in \mathbb{R}^3$
 - 4. $Q(X) < 0$ for some $X \in \mathbb{R}^3$

93. Consider the matrix

$$A(x) = \begin{pmatrix} 1+x^2 & 7 & 11 \\ 3x & 2x & 4 \\ 8x & 17 & 13 \end{pmatrix}; x \in \mathbb{R}. \text{ Then}$$

- 1. A(x) has eigenvalue 0 for some $x \in \mathbb{R}$

- 2. 0 is not an eigenvalue of A(x) for any $x \in \mathbb{R}$
- 3. A(x) has eigenvalue 0 for all $x \in \mathbb{R}$
- 4. A(x) is invertible for every $x \in \mathbb{R}$

DECEMBER - 2017

PART - B

94. Let A be a real symmetric matrix and $B = I + iA$, where $i^2 = -1$. Then
- 1. B is invertible if and only if A is invertible
 - 2. all eigenvalues of B are necessarily real
 - 3. $B - I$ is necessarily invertible
 - 4. B is necessarily invertible

95. Let $A = \begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix}$. Then the smallest positive integer n such that $A^n = I$ is
- 1. 1
 - 2. 2
 - 3. 4
 - 4. 6

96. Let $A = \begin{bmatrix} 1 & -1 & 1 \\ 1 & 1 & 1 \\ 2 & 3 & \alpha \end{bmatrix}$ and $b = \begin{bmatrix} 1 \\ 3 \\ \beta \end{bmatrix}$. Then the system $AX = b$ over the real numbers has
- 1. no solution whenever $\beta \neq 7$.
 - 2. an infinite number of solutions whenever $\alpha \neq 2$.
 - 3. an infinite number of solutions if $\alpha = 2$ and $\beta \neq 7$
 - 4. a unique solution if $\alpha \neq 2$

97. Let $A = \begin{bmatrix} 1 & 2 \\ 4 & 3 \end{bmatrix} \in M_2(\mathbb{R})$ and $\phi : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}$ be the bilinear map defined by $\phi(v, w) = v^T A w$. Choose the correct statement from below:

- 1. $\phi(v, w) = \phi(w, v)$ for all $v, w \in \mathbb{R}^2$
- 2. there exists nonzero $v \in \mathbb{R}^2$ such that $\phi(v, w) = 0$ for all $w \in \mathbb{R}^2$
- 3. there exists a 2×2 symmetric matrix B such that $\phi(v, v) = v^T B v$ for all $v \in \mathbb{R}^2$
- 4. the map $\psi : \mathbb{R}^4 \rightarrow \mathbb{R}$ defined by

$$\psi \left(\begin{bmatrix} v_1 \\ v_2 \\ w_1 \\ w_2 \end{bmatrix} \right) = \phi \left(\begin{bmatrix} v_1 \\ v_2 \end{bmatrix}, \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} \right) \text{ is linear}$$

PART – C

98. Let A be an $m \times n$ matrix with rank r . If the linear system $AX=b$ has a solution for each $b \in \mathbb{R}^m$, then
1. $m=r$
 2. the column space of A is a proper subspace of \mathbb{R}^m
 3. the null space of A is a non-trivial subspace of \mathbb{R}^n whenever $m=n$
 4. $m \geq n$ implies $m=n$

99. Let $M = \left\{ A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbb{Z} \right\}$ and the eigenvalues of A are in \mathbb{Q} . Then

1. M is empty
2. $M = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbb{Z} \right\}$
3. If $A \in M$ then the eigenvalues of A are in \mathbb{Z}
4. If $A, B \in M$ are such that $AB=1$ then $\det A \in \{+1, -1\}$

100. Let A be a 3×3 matrix with real entries. Identify the correct statements.
1. A is necessarily diagonalizable over \mathbb{R}
 2. If A has distinct real eigenvalues then it is diagonalizable over \mathbb{R}
 3. If A has distinct eigenvalues then it is diagonalizable over \mathbb{C}
 4. If all eigenvalues of A are non-zero then it is diagonalizable over \mathbb{C}

101. Let V be the vector space over \mathbb{C} of all polynomials in a variable X of degree at most 3. Let $D:V \rightarrow V$ be the linear operator given by differentiation with respect to X . Let A be the matrix of D with respect to some basis for V . Which of the following are true?
1. A is a nilpotent matrix
 2. A is a diagonalizable matrix
 3. the rank of A is 2
 4. The Jordan canonical form of A is

$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

102. For every 4×4 real symmetric non-singular matrix A , there exists a positive integer p such that
1. $pI + A$ is positive definite
 2. A^p is positive definite
 3. A^{-p} is positive definite
 4. $\exp(pA) - I$ is positive definite

103. Let A be an $m \times n$ matrix of rank m with $n > m$. If for some non-zero real number α , we have $x^t A A^t x = \alpha x^t x$, for all $x \in \mathbb{R}^m$ then $A^t A$ has
1. exactly two distinct eigenvalues
 2. 0 as an eigenvalue with multiplicity $n-m$
 3. α as a non-zero eigenvalue
 4. exactly two non-zero distinct eigenvalues

JUNE – 2018

PART – B

104. Let \mathbb{R}^n , $n \geq 2$, be equipped with standard inner product. Let $\{v_1, v_2, \dots, v_n\}$ be n column vectors forming an orthonormal basis of \mathbb{R}^n . Let A be the $n \times n$ matrix formed by the column vectors v_1, \dots, v_n . Then
1. $A = A^{-1}$
 2. $A = A^T$
 3. $A^{-1} = A^T$
 4. $\det(A) = 1$

105. Let A be a $(m \times n)$ matrix and B be a $(n \times m)$ matrix over real numbers with $m < n$. Then
1. AB is always nonsingular
 2. AB is always singular
 3. BA is always nonsingular
 4. BA is always singular

106. If A is a (2×2) matrix over \mathbb{R} with $\det(A+I) = 1 + \det(A)$, then we can conclude that
1. $\det(A) = 0$
 2. $A=0$
 3. $\text{Tr}(A) = 0$
 4. A is nonsingular

107. The system of equations:
- $$-1 \cdot x + 2 \cdot x^2 + 3 \cdot xy + 0 \cdot y = 6$$
- $$2 \cdot x + 1 \cdot x^2 + 3 \cdot xy + 1 \cdot y = 5$$
- $$3 \cdot x - 1 \cdot x^2 + 0 \cdot xy + 1 \cdot y = 7$$
1. has solutions in rational numbers
 2. has solutions in real numbers
 3. has solutions in complex numbers
 4. has no solution

108. The trace of the matrix $\begin{pmatrix} 2 & 1 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{pmatrix}^{20}$ is
1. 7^{20}
 2. $2^{20} + 3^{20}$
 3. $2 \cdot 2^{20} + 3^{20}$
 4. $2^{20} + 3^{20} + 1$

PART - C

109. Let $A = \begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & -2 \\ 0 & 0 & 1 \end{pmatrix}$ and define for $x, y,$

$$z \in \mathbb{R} \quad Q(x, y, z) = (x \ y \ z) A \begin{pmatrix} x \\ y \\ z \end{pmatrix}.$$

Which of the following statements are true?

1. The matrix of second order partial derivatives of the quadratic form Q is $2A$.
 2. The rank of the quadratic form Q is 2
 3. The signature of the quadratic form Q is $(+ + 0)$
 4. The quadratic form Q takes the value 0 for some non-zero vector (x, y, z)
110. Let $M_n(\mathbb{R})$ denote the space of all $n \times n$ real matrices identified with the Euclidean space \mathbb{R}^{n^2} . Fix a column vector $x \neq 0$ in \mathbb{R}^n . Define $f: M_n(\mathbb{R}) \rightarrow \mathbb{R}$ by $f(A) = \langle A^2x, x \rangle$. Then
1. f is linear
 2. f is differentiable
 3. f is continuous but not differentiable
 4. f is unbounded
111. Let V denote the vector space of all sequences $\mathbf{a} = (a_1, a_2, \dots)$ of real numbers such that $\sum 2^n |a_n|$ converges. Define $\|\cdot\|: V \rightarrow \mathbb{R}$ by $\|\mathbf{a}\| = \sum 2^n |a_n|$. Which of the following are true?
1. V contains only the sequence $(0, 0, \dots)$
 2. V is finite dimensional
 3. V has a countable linear basis
 4. V is a complete normed space
112. Let V be a vector space over \mathbb{C} with dimension n . Let $T: V \rightarrow V$ be a linear transformation with only 1 as eigenvalue. Then which of the following must be true?
1. $T - I = 0$
 2. $(T - I)^{n-1} = 0$
 3. $(T - I)^n = 0$
 4. $(T - I)^{2n} = 0$

113. If A is a (5×5) matrix and the dimension of the solution space of $Ax = 0$ is at least two, then

1. $\text{Rank}(A^2) \leq 3$
2. $\text{Rank}(A^2) \geq 3$
3. $\text{Rank}(A^2) = 3$
4. $\text{Det}(A^2) = 0$

114. Let $A \in M_3(\mathbb{R})$ be such that $A^8 = I_{3 \times 3}$. Then

1. minimal polynomial of A can only be of degree 2
2. minimal polynomial of A can only be of degree 3
3. either $A = I_{3 \times 3}$ or $A = -I_{3 \times 3}$
4. there are uncountably many A satisfying the above

115. Let A be an $n \times n$ matrix (with $n > 1$) satisfying $A^2 - 7A + 12I_{n \times n} = O_{n \times n}$, where $I_{n \times n}$ and $O_{n \times n}$ denote the identity matrix and zero matrix of order n respectively. Then which of the following statements are true?

1. A is invertible
2. $t^2 - 7t + 12n = 0$ where $t = \text{Tr}(A)$
3. $d^2 - 7d + 12 = 0$ where $d = \text{Det}(A)$
4. $\lambda^2 - 7\lambda + 12 = 0$ where λ is an eigenvalue of A

116. Let A be a (6×6) matrix over \mathbb{R} with characteristic polynomial $(x - 3)^2 (x - 2)^4$ and minimal polynomial $(x - 3)(x - 2)^2$. Then Jordan canonical form of A can be

1. $\begin{pmatrix} 3 & 0 & 0 & 0 & 0 & 0 \\ 0 & 3 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 1 & 0 & 0 \\ 0 & 0 & 0 & 2 & 1 & 0 \\ 0 & 0 & 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 0 & 0 & 2 \end{pmatrix}$
2. $\begin{pmatrix} 3 & 0 & 0 & 0 & 0 & 0 \\ 0 & 3 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 1 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 2 \end{pmatrix}$
3. $\begin{pmatrix} 3 & 0 & 0 & 0 & 0 & 0 \\ 0 & 3 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 1 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 0 & 0 & 2 \end{pmatrix}$

$$4. \begin{pmatrix} 3 & 1 & 0 & 0 & 0 & 0 \\ 0 & 3 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 1 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 0 & 0 & 2 \end{pmatrix}$$

117. Let V be an inner product space and S be a subset of V . Let \bar{S} denote the closure of S in V with respect to the topology induced by the metric given by the inner product. Which of the following statements are true?

1. $S = (S^\perp)^\perp$
2. $\bar{S} = (S^\perp)^\perp$
3. $\overline{\text{span}(S)} = (S^\perp)^\perp$
4. $S^\perp = ((S^\perp)^\perp)^\perp$

DECEMBER – 2018

PART – B

118. Consider the subspaces W_1 and W_2 of \mathbb{R}^3 given by $W_1 = \{(x,y,z) \in \mathbb{R}^3 : x + y + z = 0\}$ and $W_2 = \{(x,y,z) \in \mathbb{R}^3 : x - y + z = 0\}$. If W is a subspace of \mathbb{R}^3 such that

- (i) $W \cap W_2 = \text{span}\{(0,1,1)\}$
- (ii) $W \cap W_1$ is orthogonal to $W \cap W_2$ with respect to the usual inner product of \mathbb{R}^3 , then

1. $W = \text{span}\{(0,1,-1), (0,1,1)\}$
2. $W = \text{span}\{(1,0,-1), (0,1,-1)\}$
3. $W = \text{span}\{(1,0,-1), (0,1,1)\}$
4. $W = \text{span}\{(1,0,-1), (1,0,1)\}$

119. Let $C = \left[\begin{pmatrix} 1 \\ 2 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \end{pmatrix} \right]$ be a basis of \mathbb{R}^2 and $T:$

$$\mathbb{R}^2 \rightarrow \mathbb{R}^2 \text{ be defined by } T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x+y \\ x-2y \end{pmatrix}. \text{ If}$$

$T[C]$ represents the matrix of T with respect to the basis C , then which among the following is true?

1. $T[C] = \begin{bmatrix} -3 & -2 \\ 3 & 1 \end{bmatrix}$

$$2. T[C] = \begin{bmatrix} 3 & -2 \\ -3 & 1 \end{bmatrix}$$

$$3. T[C] = \begin{bmatrix} -3 & -1 \\ 3 & 2 \end{bmatrix}$$

$$4. T[C] = \begin{bmatrix} 3 & -1 \\ -3 & 2 \end{bmatrix}$$

120. Let $W_1 = \{(u,v,w,x) \in \mathbb{R}^4 \mid u+v+w=0, 2v+x=0, 2u+2w-x=0\}$ and

$W_2 = \{(u,v,w,x) \in \mathbb{R}^4 \mid u+w+x=0, u+w-2x=0, v-x=0\}$. Then which of the following is true?

1. $\dim(W_1) = 1$
2. $\dim(W_2) = 2$
3. $\dim(W_1 \cap W_2) = 1$
4. $\dim(W_1 + W_2) = 3$

121. Let A be an $n \times n$ complex matrix. Assume that A is self-adjoint and let B denotes the inverse of $(A + iI)$. Then all eigenvalues of

$(A - iI_n)B$ are

1. purely imaginary
2. of modulus one
3. real
4. of modulus less than one

122. Let $\{u_1, u_2, \dots, u_n\}$ be an orthonormal basis of \mathbb{C}^n as column vectors. Let $M = (u_1, \dots, u_k)$, $N = (u_{k+1}, \dots, u_n)$ and P be the diagonal $k \times k$ matrix with diagonal entries $\alpha_1, \alpha_2, \dots, \alpha_k \in \mathbb{R}$. Then which of the following is true?

1. $\text{Rank}(MPM^*) = k$, whenever $\alpha_i \neq \alpha_j, 1 \leq i, j \leq k$.
2. $\text{Trace}(MPM^*) = \sum_{i=1}^k \alpha_i$
3. $\text{Rank}(M^*N) = \min(k, n-k)$
4. $\text{Rank}(MM^* + NN^*) < n$

123. Let $B: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ be the function $B(a,b) = ab$. Which of the following is true?

1. B is a linear transformation
2. B is a positive definite bilinear form
3. B is symmetric but not positive definite
4. B is neither linear nor bilinear

PART – C

124. Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a linear map that satisfies $T^2 = T - I_n$. Then which of the following are true?

1. T is invertible

2. $T - I_n$ is not invertible
3. T has a real eigen value
4. $T^3 = -I_n$

125. Let $M = \begin{bmatrix} 2 & 0 & 3 & 2 & 0 & -2 \\ 0 & 1 & 0 & -1 & 3 & 4 \\ 0 & 0 & 1 & 0 & 4 & 4 \\ 1 & 1 & 1 & 0 & 1 & 1 \end{bmatrix}$,

$b_1 = \begin{bmatrix} 5 \\ 1 \\ 1 \\ 4 \end{bmatrix}$ and $b_2 = \begin{bmatrix} 5 \\ 1 \\ 3 \\ 3 \end{bmatrix}$. Then which of the

following are true?

1. both systems $MX = b_1$ and $MX = b_2$ are inconsistent
2. both systems $MX = b_1$ and $MX = b_2$ are consistent
3. the system $MX = b_1 - b_2$ is consistent
4. the systems $MX = b_1 - b_2$ is inconsistent

126. Let $M = \begin{bmatrix} 1 & -1 & 1 \\ 2 & 1 & 4 \\ -2 & 1 & -4 \end{bmatrix}$. Given that 1 is an

eigen value of M , then which among the following are correct?

1. The minimal polynomial of M is $(X - 1)(X + 4)$
2. The minimal polynomial of M is $(X - 1)^2(X + 4)$
3. M is not diagonalizable
4. $M^{-1} = \frac{1}{4}(M + 3I)$

127. Let A be a real matrix with characteristic polynomial $(X - 1)^3$. Pick the correct statements from below:

1. A is necessarily diagonalizable
2. If the minimal polynomial of A is $(X - 1)^3$, then A is diagonalizable
3. Characteristic polynomial of A^2 is $(X - 1)^3$
4. If A has exactly two Jordan blocks, then $(A - I)^2$ is diagonalizable

128. Let P_3 be the vector space of polynomials with real coefficients and of degree at most 3. Consider the linear map $T : P_3 \rightarrow P_3$ defined by $T(p(x)) = p(x + 1) + p(x - 1)$. Which of the following properties does the matrix of T (with respect to the standard basis $B = \{1, x, x^2, x^3\}$ of P_3) satisfy?

1. $\det T = 0$
2. $(T - 2I)^4 = 0$ but $(T - 2I)^3 \neq 0$

3. $(T - 2I)^3 = 0$ but $(T - 2I)^2 \neq 0$
4. 2 is an eigen value with multiplicity 4

129. Let M be an $n \times n$ Hermitian matrix of rank k , $k \neq n$. If $\lambda \neq 0$ is an eigen value of M with corresponding unit column vector u , with $Mu = \lambda u$, then which of the following are true?

1. $\text{rank}(M - \lambda uu^*) = k - 1$
2. $\text{rank}(M - \lambda uu^*) = k$
3. $\text{rank}(M - \lambda uu^*) = k + 1$
4. $(M - \lambda uu^*)^n = M^n - \lambda^n uu^*$

130. Define a real valued function B on $\mathbb{R}^2 \times \mathbb{R}^2$ as follows. If $u = (x_1, x_2)$, $w = (y_1, y_2)$ belong to \mathbb{R}^2 define $B(u, w) = x_1y_1 - x_1y_2 - x_2y_1 + 4x_2y_2$. Let $v_0 = (1, 0)$ and let $W = \{v \in \mathbb{R}^2 : B(v_0, v) = 0\}$. Then W

1. is not a subspace of \mathbb{R}^2
2. equals $\{(0, 0)\}$
3. is the y axis
4. is the line passing through $(0, 0)$ and $(1, 1)$

131. Consider the Quadratic forms

$$Q_1(x, y) = xy$$

$$Q_2(x, y) = x^2 + 2xy + y^2$$

$Q_3(x, y) = x^2 + 3xy + 2y^2$ on \mathbb{R}^2 . Choose the correct statements from below:

1. Q_1 and Q_2 are equivalent
2. Q_1 and Q_3 are equivalent
3. Q_2 and Q_3 are equivalent
4. all are equivalent

JUNE-2019

PART - B

132. Consider the vector space P_n of real polynomials in x of degree less than or equal to n . Define $T : P_2 \rightarrow P_3$ by $(Tf)(x) = \int_0^x f(t) dt + f'(x)$. Then the matrix representation of T with respect to the bases $\{1, x, x^2\}$ and $\{1, x, x^2, x^3\}$ is

1. $\begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & \frac{1}{2} & 0 \\ 0 & 2 & 0 & \frac{1}{3} \end{pmatrix}$ 2. $\begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 2 \\ 0 & \frac{1}{2} & 0 \\ 0 & 0 & \frac{1}{3} \end{pmatrix}$

3.
$$\begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 2 & 0 \\ 0 & \frac{1}{2} & 0 & \frac{1}{3} \end{pmatrix}$$

4.
$$\begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & \frac{1}{2} \\ 0 & 2 & 0 \\ 0 & 0 & \frac{1}{3} \end{pmatrix}$$

- 133.** Let $P_A(x)$ denote the characteristic polynomial of a matrix A . Then for which of the following matrices, $P_A(x) - P_{A^{-1}}(x)$ is a constant?

1.
$$\begin{pmatrix} 3 & 3 \\ 2 & 4 \end{pmatrix}$$

2.
$$\begin{pmatrix} 4 & 3 \\ 2 & 3 \end{pmatrix}$$

3.
$$\begin{pmatrix} 3 & 2 \\ 4 & 3 \end{pmatrix}$$

4.
$$\begin{pmatrix} 2 & 3 \\ 3 & 4 \end{pmatrix}$$

- 134.** Which of the following matrices is not diagonalizable over \mathbb{R} ?

1.
$$\begin{pmatrix} 2 & 0 & 1 \\ 0 & 3 & 0 \\ 0 & 0 & 2 \end{pmatrix}$$

2.
$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$$

3.
$$\begin{pmatrix} 2 & 1 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{pmatrix}$$

4.
$$\begin{pmatrix} 1 & -1 \\ 2 & 4 \end{pmatrix}$$

- 135.** What is the rank of the following matrix?

$$\begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ 1 & 2 & 2 & 2 & 2 \\ 1 & 2 & 3 & 3 & 3 \\ 1 & 2 & 3 & 4 & 4 \\ 1 & 2 & 3 & 4 & 5 \end{pmatrix}$$

1. 2
3. 4

2. 3
4. 5

- 136.** Let V denote the vector space of real valued continuous functions on the closed interval $[0, 1]$. Let W be the subspace of V spanned by $\{\sin(x), \cos(x), \tan(x)\}$. Then the dimension of W over \mathbb{R} is

1. 1
3. 3
2. 2
4. infinite

- 137.** Let V be the vector space of polynomials in the variable t of degree at most 2 over \mathbb{R} . An inner product on V is defined by

$$\langle f, g \rangle = \int_0^1 f(t)g(t)dt$$

for $f, g \in V$. Let $W = \text{span}\{1 - t^2, 1 + t^2\}$ and W^\perp be the orthogonal complement of W in V . Which of the following conditions is satisfied for all $h \in W^\perp$?

1. h is an even function, i.e. $h(t) = h(-t)$
2. h is an odd function, i.e. $h(t) = -h(-t)$
3. $h(t) = 0$ has a real solution
4. $h(0) = 0$

PART - C

- 138.** Let $L(\mathbb{R}^n)$ be the space of \mathbb{R} -linear maps from \mathbb{R}^n to \mathbb{R}^n . If $\text{Ker}(T)$ denotes the kernel (null space) of T then which of the following are true?

1. There exists $T \in L(\mathbb{R}^5) \setminus \{0\}$ such that $\text{Range}(T) = \text{Ker}(T)$
2. There does not exist $T \in L(\mathbb{R}^5) \setminus \{0\}$ such that $\text{Range}(T) = \text{Ker}(T)$
3. There exists $T \in L(\mathbb{R}^6) \setminus \{0\}$ such that $\text{Range}(T) = \text{Ker}(T)$
4. There does not exist $T \in L(\mathbb{R}^6) \setminus \{0\}$ such that $\text{Range}(T) = \text{Ker}(T)$

- 139.** Let V be a finite dimensional vector space over \mathbb{R} and $T : V \rightarrow V$ be a linear map. Can you always write $T = T_2 \circ T_1$ for some linear maps $T_1 : V \rightarrow W, T_2 : W \rightarrow V$, where W is some finite dimensional vector space and such that

1. both T_1 and T_2 are onto
2. both T_1 and T_2 are one to one
3. T_1 is onto, T_2 is one to one
4. T_1 is one to one, T_2 is onto

- 140.** Let $A = ((a_{ij}))$ be a 3×3 complex matrix. Identify the correct statements

1. $\det(((-1)^{i+j} a_{ij})) = \det A$
2. $\det(((-1)^{i+j} a_{ij})) = -\det A$
3. $\det(((\sqrt{-1})^{i+j} a_{ij})) = \det A$
4. $\det(((\sqrt{-1})^{i+j} a_{ij})) = -\det A$

- 141.** Let $p(x) = a_0 + a_1x + \dots + a_nx^n$ be a non-constant polynomial of degree $n \geq 1$. Consider the polynomial

$$q(x) = \int_0^x p(t)dt, r(x) = \frac{d}{dx} p(x).$$

Let V denote the real vector space of all polynomials in x . Then which of the following are true?

- 1. q and r are linearly independent in V
- 2. q and r are linearly dependent in V
- 3. x^n belongs to the linear span of q and r
- 4. x^{n+1} belongs to the linear span of q and r

142. Let $M_n(\mathbb{R})$ be the ring of $n \times n$ matrices over \mathbb{R} . Which of the following are true for every $n \geq 2$?

- 1. there exist matrices $A, B \in M_n(\mathbb{R})$ such that $AB - BA = I_n$, where I_n denotes the identity $n \times n$ matrix.
- 2. if $A, B \in M_n(\mathbb{R})$ and $AB = BA$, then A is diagonalizable over \mathbb{R} if and only if B is diagonalizable over \mathbb{R}
- 3. if $A, B \in M_n(\mathbb{R})$, then AB and BA have same minimal polynomial
- 4. if $A, B \in M_n(\mathbb{R})$, then AB and BA have the same eigen values in \mathbb{R}

143. Consider a matrix $A = (a_{ij})_{5 \times 5}$, $1 \leq i, j \leq 5$ such that $a_{ij} = \frac{1}{n_i + n_j + 1}$, where $n_i, n_j \in \mathbb{N}$.

Then in which of the following cases A is a positive definite matrix?

- 1. $n_i = i$ for all $i = 1, 2, 3, 4, 5$
- 2. $n_1 < n_2 < \dots < n_5$
- 3. $n_1 = n_2 = \dots = n_5$
- 4. $n_1 > n_2 > \dots > n_5$

144. Let $\langle \cdot, \cdot \rangle : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ denote the standard inner product on \mathbb{R}^n . For a non zero $w \in \mathbb{R}^n$, define $T_w : \mathbb{R}^n \rightarrow \mathbb{R}^n$ by

$$T_w(v) = v - \frac{2\langle v, w \rangle}{\langle w, w \rangle} w, \text{ for } v \in \mathbb{R}^n. \text{ Which of}$$

the following are true?

- 1. $\det(T_w) = 1$
- 2. $\langle T_w(v_1), T_w(v_2) \rangle = \langle v_1, v_2 \rangle \forall v_1, v_2 \in \mathbb{R}^n$
- 3. $T_w = T_w^{-1}$
- 4. $T_{2w} = 2T_w$

145. Consider the matrix $A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ over the

field \mathbb{Q} of rationals. Which of the following matrices are of the form $P^t AP$ for a suitable 2×2 invertible matrix P over \mathbb{Q} ? Here P^t denotes the transpose of P.

- 1. $\begin{pmatrix} 2 & 0 \\ 0 & -2 \end{pmatrix}$
- 2. $\begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$

- 3. $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$
- 4. $\begin{pmatrix} 3 & 4 \\ 4 & 5 \end{pmatrix}$

DECEMBER - 2019

PART - B

146. Let $A = \begin{pmatrix} 2 & 0 & 5 \\ 1 & 2 & 3 \\ -1 & 5 & 1 \end{pmatrix}$. The system of linear

equations $AX = Y$ has a solution

1. only for $Y = \begin{pmatrix} x \\ 0 \\ 0 \end{pmatrix}, x \in \mathbb{R}$

2. only for $Y = \begin{pmatrix} 0 \\ y \\ 0 \end{pmatrix}, y \in \mathbb{R}$

3. only for $Y = \begin{pmatrix} 0 \\ y \\ z \end{pmatrix}, y, z \in \mathbb{R}$

4. for all $Y \in \mathbb{R}^3$

147. Let V be a vector space of dimension 3 over \mathbb{R} . Let $T : V \rightarrow V$ be a linear transformation,

given by the matrix $A = \begin{pmatrix} 1 & -1 & 0 \\ 1 & -4 & 3 \\ -2 & 5 & -3 \end{pmatrix}$ with

respect to an ordered basis (v_1, v_2, v_3) of V. Then which of the following statements is true?

- 1. $T(v_3) = 0$
- 2. $T(v_1 + v_2) = 0$
- 3. $T(v_1 + v_2 + v_3) = 0$
- 4. $T(v_1 + v_3) = T(v_2)$

148. Let $M_4(\mathbb{R})$ be the space of all (4×4) matrices over \mathbb{R} . Let

$$W = \left\{ (a_{ij}) \in M_4(\mathbb{R}) \mid \sum_{i+j=k} a_{ij} = 0, \text{ for } k = 2, 3, 4, 5, 6, 7, 8 \right\}$$

Then $\dim(W)$ is

- 1. 7
- 2. 8
- 3. 9
- 4. 10

149. For $t \in \mathbb{R}$, define

$$M(t) = \begin{pmatrix} 1 & t & 0 \\ 1 & 1 & t^2 \\ 0 & 1 & 1 \end{pmatrix}.$$

Then which of the following statements is true?

1. $\det M(t)$ is a polynomial function of degree 3 in t
 2. $\det M(t) = 0$ for all $t \in \mathbb{R}$
 3. $\det M(t)$ is zero for infinitely many $t \in \mathbb{R}$
 4. $\det M(t)$ is zero for exactly two $t \in \mathbb{R}$
150. For a quadratic form in 3 variables over \mathbb{R} , let r be the rank and s be the signature. The number of possible pairs (r, s) is
1. 13
 2. 9
 3. 10
 4. 16

PART - C

151. Let $A \in M_3(\mathbb{R})$ and let $X = \{C \in GL_3(\mathbb{R}) \mid CAC^{-1} \text{ is triangular}\}$. Then
1. $X \neq \emptyset$
 2. If $X = \emptyset$, then A is not diagonalizable over \mathbb{C}
 3. If $X = \emptyset$. Then A is diagonalizable over \mathbb{C}
 4. If $X = \emptyset$, then A has no real eigenvalue
152. Which of the following statements regarding quadratic forms in 3 variables are true?
1. Any two quadratic forms of rank 3 are isomorphic over \mathbb{R}
 2. Any two quadratic forms of rank 3 are isomorphic over \mathbb{C}
 3. There are exactly three non zero quadratic forms of rank ≤ 3 upto isomorphism over \mathbb{C}
 4. There are exactly three non zero quadratic forms of rank 2 upto isomorphism over \mathbb{R} and \mathbb{C}
153. Let $T : \mathbb{C}^n \rightarrow \mathbb{C}^n$ be a linear transformation, $n \geq 2$. Suppose 1 is the only eigenvalue of T . Which of the following statements are true?
1. $T^k \neq I$ for any $k \in \mathbb{N}$
 2. $(T - I)^{n-1} = 0$
 3. $(T - I)^n = 0$
 4. $(T - I)^{n+1} = 0$

154. Let X be a finite dimensional inner product space over \mathbb{C} . Let $T : X \rightarrow X$ be any linear

transformation. Then which of the following statements are true?

1. T is unitary $\Rightarrow T$ is self adjoint
2. T is self adjoint $\Rightarrow T$ is normal
3. T is unitary $\Rightarrow T$ is normal
4. T is normal $\Rightarrow T$ is unitary

155. Let $n \geq 1$ and $\alpha, \beta \in \mathbb{R}$ with $\alpha \neq \beta$. Suppose $A_n(\alpha, \beta) = [a_{ij}]$ is an $n \times n$ matrix such that $a_{ij} = \alpha$ and $a_{ij} = \beta$ for $i \neq j$, $1 \leq i, j \leq n$. Let D_n be the determinant of $A_n(\alpha, \beta)$. Which of the following statements are true?

1. $D_n = (\alpha - \beta)D_{n-1} + \beta$ for $n \geq 2$
2. $\frac{D_n}{(\alpha - \beta)^{n-1}} = \frac{D_{n-1}}{(\alpha - \beta)^{n-2}} + \beta$ for $n \geq 2$
3. $D_n = (\alpha + (n - 1)\beta)^{n-1}(\alpha - \beta)$ for $n \geq 2$
4. $D_n = (\alpha + (n - 1)\beta)(\alpha - \beta)^{n-1}$ for $n \geq 2$

156. Which of the following statements are true?

1. Any two quadratic forms of same rank in n -variables over \mathbb{R} are isomorphic
2. Any two quadratic forms of same rank in n -variables over \mathbb{C} are isomorphic
3. Any two quadratic forms in n -variables are isomorphic over \mathbb{C}
4. A quadratic form in 4 variables may be isomorphic to a quadratic form in 10 variables

157. Let $T : \mathbb{R}^4 \rightarrow \mathbb{R}^4$ be a linear transformation with characteristic polynomial $(x - 2)^4$ and minimal polynomial $(x - 2)^2$. Jordan canonical form of T can be

1. $\begin{pmatrix} 2 & 0 & 0 & 0 \\ 1 & 2 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 1 & 2 \end{pmatrix}$ 2. $\begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 1 & 2 \end{pmatrix}$

3. $\begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 2 \end{pmatrix}$ 4. $\begin{pmatrix} 2 & 0 & 0 & 0 \\ 1 & 2 & 0 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 2 \end{pmatrix}$

JUNE-2020

PART - B

158. Let A be an $n \times n$ matrix such that the set of all its non-zero eigenvalues has exactly r elements. Which of the following statements is true?

- 1. rank $A \leq r$
- 2. If $r = 0$, then rank $A < n - 1$
- 3. rank $A \geq r$
- 4. A^2 has r distinct non zero eigenvalues

159. Let A and B be 2×2 matrices. Then which of the following is true?
- 1. $\det(A + B) + \det(A - B) = \det A + \det B$
 - 2. $\det(A + B) + \det(A - B) = 2\det A - 2\det B$
 - 3. $\det(A + B) + \det(A - B) = 2\det A + 2\det B$
 - 4. $\det(A + B) - \det(A - B) = 2\det A - 2\det B$

160. If $A = \begin{pmatrix} 3 & -2 \\ 2 & -1 \end{pmatrix}$, then A^{20} equals

- 1. $\begin{pmatrix} 41 & 40 \\ -40 & -39 \end{pmatrix}$
- 2. $\begin{pmatrix} 41 & -40 \\ 40 & -39 \end{pmatrix}$
- 3. $\begin{pmatrix} 41 & -40 \\ -40 & -39 \end{pmatrix}$
- 4. $\begin{pmatrix} 41 & 40 \\ 40 & -39 \end{pmatrix}$

161. Let A be a 2×2 real matrix with $\det A = 1$ and trace $A = 3$. What is the value of trace A^2 ?
- 1. 2
 - 2. 10
 - 3. 9
 - 4. 7

162. For $a, b \in \mathbb{R}$, let $p(x, y) = a^2x_1y_1 + abx_2y_1 + abx_1y_2 + b^2x_2y_2$, $x = (x_1, x_2), y = (y_1, y_2) \in \mathbb{R}^2$. For what values of a and b does $p : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}$ define an inner product?
- 1. $a > 0, b > 0$
 - 2. $ab > 0$
 - 3. $a = 0, b = 0$
 - 4. For no values of a, b

163. Which of the following real quadratic forms on \mathbb{R}^2 is positive definite?
- 1. $Q(X, Y) = XY$
 - 2. $Q(X, Y) = X^2 - XY + Y^2$
 - 3. $Q(X, Y) = X^2 + 2XY + Y^2$
 - 4. $Q(X, Y) = X^2 + XY$

PART - C

164. Let P be a square matrix such that $P^2 = P$. Which of the following statements are true?
- 1. Trace of P is an irrational number
 - 2. Trace of $P =$ rank of P
 - 3. Trace of P is an integer
 - 4. Trace of P is an imaginary complex number

165. Let A and B be $n \times n$ real matrices and let $C = \begin{pmatrix} A & B \\ B & A \end{pmatrix}$.

Which of the following statements are true?

- 1. If λ is an eigenvalue of $A + B$ then λ is an eigenvalue of C
- 2. If λ is an eigenvalue of $A - B$ then λ is an eigenvalue of C
- 3. If λ is an eigenvalue of A or B then λ is an eigenvalue of C
- 4. All eigenvalues of C are real

166. Let A be an $n \times n$ real matrix. Let b be an $n \times 1$ vector. Suppose $Ax = b$ has no solution. Which of the following statements are true?
- 1. There exists an $n \times 1$ vector c such that $Ax = c$ has a unique solution
 - 2. There exist infinitely many vectors c such that $Ax = c$ has no solution
 - 3. If y is the first column of A then $Ax = y$ has a unique solution
 - 4. $\det A = 0$

167. Let A be an $n \times n$ matrix such that the first 3 rows of A are linearly independent and the first 5 columns of A are linearly independent. Which of the following statements are true?
- 1. A has atleast 5 linearly independent rows
 - 2. $3 \leq \text{rank } A \leq 5$
 - 3. $\text{rank } A \geq 5$
 - 4. $\text{rank } A^2 \geq 5$

168. Let n be a positive integer and F be a non-empty proper subset of $\{1, 2, \dots, n\}$. Define $\langle x, y \rangle_F = \sum_{k \in F} x_k y_k$, $x = (x_1, \dots, x_n), y = (y_1, \dots, y_n) \in \mathbb{R}^n$.

Let $T = \{x \in \mathbb{R}^n : \langle x, x \rangle_F = 0\}$. Which of the following statements are true?

For $y \in \mathbb{R}^n, y \neq 0$

- 1. $\inf_{x \in T} \langle x + y, x + y \rangle_F = \langle y, y \rangle_F$
- 2. $\sup_{x \in T} \langle x + y, x + y \rangle_F = \langle y, y \rangle_F$
- 3. $\inf_{x \in T} \langle x + y, x + y \rangle_F < \langle y, y \rangle_F$
- 4. $\sup_{x \in T} \langle x + y, x + y \rangle_F > \langle y, y \rangle_F$

169. Let $v \in \mathbb{R}^3$ be a non-zero vector. Define a linear transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ by

$$T(x) = x - 2 \frac{x \cdot v}{v \cdot v} v, \text{ where } x \cdot y \text{ denotes the}$$

standard inner product in \mathbb{R}^3 . Which of the following statements are true?

1. The eigenvalues of T are +1, -1
2. The determinant of T is -1
3. The trace of T is +1
4. T is distance preserving

170. A quadratic form $Q(x,y,z)$ over \mathbb{R} represents

0 non trivially if there exists $(a,b,c) \in \mathbb{R}^3 \setminus \{(0,0,0)\}$ such that $Q(a, b, c) = 0$. Which of the following quadratic forms $Q(x, y, z)$ over

\mathbb{R} represent 0 non trivially?

1. $Q(x, y, z) = xy + z^2$
2. $Q(x, y, z) = x^2 + 3y^2 - 2z^2$
3. $Q(x, y, z) = x^2 - xy + y^2 + z^2$
4. $Q(x, y, z) = x^2 + xy + z^2$

171. Let $Q(x, y, z)$ be a real quadratic form. Which of the following statements are true?

1. $Q(x_1 + x_2, y, z) = Q(x_1, y, z) + Q(x_2, y, z)$ for all x_1, x_2, y, z
2. $Q(x_1 + x_2, y_1 + y_2, 0) + Q(x_1 - x_2, y_1 - y_2, 0) = 2Q(x_1, y_1, 0) + 2Q(x_2, y_2, 0)$ for all x_1, x_2, y_1, y_2
3. $Q(x_1 + x_2, y_1 + y_2, z_1 + z_2) = Q(x_1, y_1, z_1) + Q(x_2, y_2, z_2)$ for at least one choice of $x_1, x_2, y_1, y_2, z_1, z_2$
4. $2Q(x_1 + x_2, y_1 + y_2, 0) + 2Q(x_1 - x_2, y_1 - y_2, 0) = Q(x_1, y_1, 0) + Q(x_2, y_2, 0)$ for all x_1, x_2, y_1, y_2

JUNE-2021

PART - B

172. Let A be a 4×4 matrix such that -1, 1, 1, -2 are its eigenvalues. If $B = A^4 - 5A^2 + 5I$, then trace (A + B) equals

- (1) 0
- (2) -12
- (3) 3
- (4) 9

173. Let $M = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 2 & -1 \\ 1 & 1 & 3 \end{bmatrix}$. Given that 1 is an

eigenvalue of M, which of the following statements is true?

- (1) -2 is an eigenvalue of M
- (2) 3 is an eigenvalue of M
- (3) The eigen space of each eigen value has dimension 1
- (4) M is diagonalizable

174. Let A and B be $n \times n$ matrices. Suppose the sum of the elements in any row of A is 2 and the sum of the elements in any column of B is 2. Which of the following matrices is necessarily singular?

- (1) $I - \frac{1}{2} BA^T$
- (2) $I - \frac{1}{2} AB$
- (3) $I - \frac{1}{4} AB$
- (4) $I - \frac{1}{4} BA^T$

175. Let $V = \{A \in M_{3 \times 3}(\mathbb{R}) : A^t + A \in \mathbb{R} \cdot I\}$ where I is the identity matrix. Consider the quadratic form defined as $q(A) = \text{Trace}(A)^2 - \text{Trace}(A^2)$. What is the signature of the quadratic form?

- (1) (+ + + +)
- (2) (+ 0 0 0)
- (3) (+ - - -)
- (4) (- - - 0)

176. Let $n > 1$ be a fixed natural number. Which of the following is an inner product on the vector space of $n \times n$ real symmetric matrices?

- (1) $\langle A, B \rangle = (\text{trace}(A)) (\text{trace}(B))$
- (2) $\langle A, B \rangle = \text{trace}(AB)$
- (3) $\langle A, B \rangle = \text{determinant}(AB)$
- (4) $\langle A, B \rangle = \text{trace}(A) + \text{trace}(B)$

177. Consider the two statements given below:

- I. There exists a matrix $N \in M_4(\mathbb{R})$ such that $\{(1, 1, 1, -1), (1, -1, 1, 1)\}$ is a basis of $\text{Row}(N)$ and $(1, 2, 1, 4) \in \text{Null}(N)$
- II. There exists a matrix $M \in M_4(\mathbb{R})$ such that $\{(1, 1, 1, 0)^T, (1, 0, 1, 1)^T\}$ is a basis of $\text{Col}(M)$ and $(1, 1, 1, 1)^T, (1, 0, 1, 0)^T \in \text{Null}(M)$

Which of the following statements is true?

- (1) Statement I is False and Statement II is True
- (2) Statement I is True and Statement II is False
- (3) Both Statement I and Statement II are False
- (4) Both Statement I and Statement II are True

PART - C

178. Let $M \in M_n(\mathbb{R})$ such that $M \neq 0$ but $M^2 = 0$. Which of the following statements are true?

- (1) If n is even then $\dim(\text{Col}(M)) > \dim(\text{Null}(M))$
- (2) If n is even then $\dim(\text{Col}(M)) \leq \dim(\text{Null}(M))$

- (3) If n is odd then $\dim(\text{Col}(M)) < \dim(\text{Null}(M))$
- (4) If n is odd then $\dim(\text{Col}(M)) > \dim(\text{Null}(M))$

179. Consider the system

$$\begin{aligned} 2x + ky &= 2 - k \\ kx + 2y &= k \\ ky + kz &= k - 1 \end{aligned}$$

in three unknowns and one real parameter k . For which of the following values of k is the system of linear equations consistent?

- (1) 1
- (2) 2
- (3) -1
- (4) -2

180. Which of the following are inner products on \mathbb{R}^2 ?

$$(1) \left\langle \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \right\rangle = x_1y_1 + 2x_1y_2 + 2x_2y_1 + x_2y_2$$

$$(2) \left\langle \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \right\rangle = x_1y_1 + x_1y_2 + x_2y_1 + 2x_2y_2$$

$$(3) \left\langle \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \right\rangle = x_1y_1 + x_1y_2 + x_2y_1 + x_2y_2$$

$$(4) \left\langle \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \right\rangle = x_1y_1 - \frac{1}{2}x_1y_2 - \frac{1}{2}x_2y_1 + x_2y_2$$

181. Let A be an $m \times n$ matrix such that the first r rows of A are linearly independent and the first s columns of A are linearly independent, where $r < m$ and $s < n$. Which of the following statements are true?

- (1) The rank of A is atleast $\max\{r, s\}$
- (2) The submatrix formed by the first r rows and the first s columns of A has rank $\min\{r, s\}$
- (3) If $r < s$, then there exists a row among rows $r + 1, \dots, m$ which together with the first r rows form a linearly independent set
- (4) If $s < r$, then there exists a column among columns $s + 1, \dots, n$ which together with the first s columns form a linearly dependent set.

182. Let A be an $n \times n$ matrix. We say that A is diagonalizable if there exists a nonsingular matrix P such that PAP^{-1} is a diagonal matrix. Which of the following conditions imply that A is diagonalizable?

- (1) There exists integer k such that $A^k = I$
- (2) There exists integer k such that A^k is nilpotent
- (3) A^2 is diagonalizable

(4) A has n linearly independent eigenvectors

183. It is known that $X = X_0 \in M_2(\mathbb{Z})$ is a solution of $AX - XA = A$ for some

$$A \in \left\{ \begin{pmatrix} 1 & 1 \\ -1 & -1 \end{pmatrix}, \begin{pmatrix} -1 & 1 \\ 1 & -1 \end{pmatrix}, \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \right\}$$

Which of the following values are NOT possible for the determinant of X_0 ?

- (1) $\det(X_0) = 0$
- (2) $\det(X_0) = 2$
- (3) $\det(X_0) = 6$
- (4) $\det(X_0) = 10$

184. Let A be an $m \times m$ matrix with real entries and let x be an $m \times 1$ vector of unknowns. Now consider the two statements given below:

I: There exists non-zero vector $b_1 \in \mathbb{R}^m$ such that the linear system $Ax = b_1$ has NO solution

II: There exist non-zero vectors $b_2, b_3 \in \mathbb{R}^m$, with $b_2 \neq cb_3$ for any $c \in \mathbb{R}$, such that the linear systems $Ax = b_2$ and $Ax = b_3$ have solutions. Which of the following statements are true?

- (1) II is TRUE whenever A is singular
- (2) I is TRUE whenever A is singular
- (3) Both I and II can be TRUE simultaneously
- (4) If $m = 2$, then at least one of I and II is FALSE

JUNE-2022

PART - B

185. Let $A = (a_{ij})$ be a real symmetric 3×3 matrix. Consider the quadratic form $Q(x_1, x_2, x_3) = x^tAx$ where $x = (x_1, x_2, x_3)^t$. Which of the following is true?

- (1) If $Q(x_1, x_2, x_3)$ is positive definite, then $a_{ij} > 0$ for all $i \neq j$.
- (2) If $Q(x_1, x_2, x_3)$ is positive definite, then $a_{i,i} > 0$ for all i .
- (3) If $a_{ij} > 0$ for all $i \neq j$, then $Q(x_1, x_2, x_3)$ is positive definite.
- (4) If $a_{i,i} > 0$ for all i , then $Q(x_1, x_2, x_3)$ is positive definite.

186. Let \mathbb{R} be the field of real numbers. Let V be the vector space of real polynomials of degree at most 1. Consider the bilinear form

$$\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R},$$

$$\text{given by } \langle f, g \rangle = \int_0^1 f(x)g(x)dx.$$

Which of the following is true?

- (1) For all non-zero real numbers a, b, there exists a real number c such that the vectors $ax + b, x + c \in V$ are orthogonal to each other.
- (2) For all non-zero real numbers b, there are infinitely many real numbers c such that the vectors $x + b, x + c \in V$ are orthogonal to each other.
- (3) For all positive real numbers c, there exist infinitely many real numbers a, b such that the vectors $ax + b, x + c \in V$ are orthogonal to each other.
- (4) For all non-zero real numbers b, there are infinitely many real numbers c such that the vectors $b, x + c \in V$ are orthogonal to each other.

187. Suppose A is a real $n \times n$ matrix of rank r. Let V be the vector space of all real $n \times n$ matrices X such that $AX = 0$. What is the dimension of V?

- (1) r (2) nr (3) n^2r (4) $n^2 - nr$

188. Suppose A and B are similar real matrices, that is, there exists an invertible matrix S such that $A = SBS^{-1}$. Which of the following need not be true?

- (1) Transpose of A is similar to the transpose of B
- (2) The minimal polynomial of A is same as the minimal polynomial of B
- (3) $\text{trace}(A) = \text{trace}(B)$
- (4) The range of A is same as the range of B

189. Let A be an invertible 5×5 matrix over a field F. Suppose that characteristic polynomials of A and A^{-1} are the same. Which of the following is necessarily true?

- (1) $\det(A)^2 = 1$ (2) $\det(A)^5 = 1$
- (3) $\text{trace}(A)^2 = 1$ (4) $\text{trace}(A)^5 = 1$

PART - C

190. Let W be the space of \mathbb{C} -linear combinations of the following functions $f_1(z) = \sin z, f_2(z) = \cos z, f_3(z) = \sin(2z), f_4(z) = \cos(2z)$

Let T be the linear operator on W given by complex differentiation. Which of the following statements are true?

- (1) Dimension of W is 3
- (2) The span of f_1 and f_2 is Jordan block of T

- (3) T has two Jordan blocks
- (4) T has four Jordan blocks

191. Let V be vector space of polynomials $f(X, Y) \in \mathbb{R}[X, Y]$ with (total) degree less than 3. Let $T : V \rightarrow V$ be the linear transformation given by $\frac{\partial}{\partial x}$. Which of the

following statements are true?

- (1) The nullity of T is atleast 3
- (2) The rank of T is atleast 4
- (3) The rank of T is atleast 3
- (4) T is invertible

192. For a positive integer $n \geq 2$, let A be an $n \times n$ matrix with entries in \mathbb{R} such that A^{n^2} has rank zero. Let 0_n denote the $n \times n$ matrix with all entries equal to 0. Which of the following statements are equivalent to the statement that A has n linearly independent eigenvectors?

- (1) $A^n = 0_n$ (2) $A^{n^2} = 0_n$
- (3) $A = 0_n$ (4) $A^2 = 0_n$

193. Let P_n be the vector space of real polynomials with degree at most n. Let \langle , \rangle be an inner product on P_n with respect to which $\left\{1, x, \frac{1}{2!}x^2, \dots, \frac{1}{n!}x^n\right\}$ is an orthonormal

basis of P_n .

Let $f = \sum_i \alpha_i x^i, g = \sum_i \beta_i x^i \in P_n$. Which of the following statements are true?

- (1) $\langle f, g \rangle = \sum_i (i!) \alpha_i \beta_i$ defines one such inner product, but there is another such inner product.
- (2) $\langle f, g \rangle = \sum_i (i!) \alpha_i \beta_i$.
- (3) $\langle f, g \rangle = \sum_i (i!)^2 \alpha_i \beta_i$ defines one such inner product, but there is another such inner product.
- (4) $\langle f, g \rangle = \sum_i (i!)^2 \alpha_i \beta_i$

194. Let U and V be the subspaces of \mathbb{R}^3 defined by

$$U = \left\{ \begin{pmatrix} x \\ y \\ z \end{pmatrix} \in \mathbb{R}^3 \mid 2x + 3y + 4z = 0 \right\}.$$

$$V = \left\{ \begin{pmatrix} x \\ y \\ z \end{pmatrix} \in \mathbb{R}^3 \mid x + 2y + 5z = 0 \right\}.$$

Which of the following statements are true?

- (1) There exists an invertible linear transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ such that $T(U) = V$.
- (2) There does not exist an invertible linear transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ such that $T(V) = U$.
- (3) There exists a linear transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ such that $T(U) \cap V \neq \{0\}$ and the characteristic polynomial of T is not the product of linear polynomials with real coefficients.
- (4) There exists a linear transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ such that $T(U) = V$ and the characteristic polynomial of T vanishes at 1.

195. Let A be an $n \times n$ matrix with entries in \mathbb{R} that A and A^2 are of same rank. Consider linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ defined by $T(v) = A(v)$ for all $v \in \mathbb{R}^n$. Which of the following statements are true.

- (1) The Kernels of T and $T \circ T$ are the same
- (2) The Kernels of T and $T \circ T$ are of equal dimensions
- (3) A must be invertible
- (4) $I_n + A$ must be invertible, where I_n denotes an $n \times n$ identity matrix

196. On the complex vector space \mathbb{C}^{100} with standard basis $\{e_1, e_2, \dots, e_{100}\}$, consider the bilinear form $B(x, y) = \sum_i x_i y_i$, where x_i and y_i are the coefficients of e_i in x and y respectively. Which of the following statements are true?

- (1) B is non-degenerate
- (2) Restriction of B to all non-zero subspaces is non-degenerate
- (3) There is a 51 dimensional subspace W of \mathbb{C}^{100} such that the restriction $B : W \times W \rightarrow \mathbb{C}$ is the zero map
- (4) There is a 49 dimensional subspace W of \mathbb{C}^{100} such that the restriction $B : W \times W \rightarrow \mathbb{C}$ is the zero map

197. For a positive integer $n \geq 2$, let $M_n(\mathbb{R})$ denote the vector space of $n \times n$ matrices with entries in \mathbb{R} . Which of the following statements are true?

- (1) The vector space $M_n(\mathbb{R})$ can be expressed as the union of a finite collection of its proper subspaces.
- (2) Let A be an element of $M_n(\mathbb{R})$. Then, for any real number x and $\varepsilon > 0$, there exists a real number $y \in (x - \varepsilon, x + \varepsilon)$ such that $\det(yI + A) \neq 0$.
- (3) Suppose A and B are two elements of $M_n(\mathbb{R})$ such that their characteristic polynomials are equal. If $A = C^2$ for some $C \in M_n(\mathbb{R})$, then $B = D^2$ for some $D \in M_n(\mathbb{R})$.
- (4) For any subspace W of $M_n(\mathbb{R})$, there exists a linear transformation $T : M_n(\mathbb{R}) \rightarrow M_n(\mathbb{R})$ with W as its image.

JUNE-2023

PART - B

198. Let T be a linear operator on \mathbb{R}^3 . Let $f(X) \in \mathbb{R}[X]$ denote its characteristic polynomial. Consider the following statements.

- (a) Suppose T is non-zero and 0 is an eigen value of T . If we write $f(X) = Xg(X)$ in $\mathbb{R}[X]$, then the linear operator $g(T)$ is zero.
- (b) Suppose 0 is an eigenvalue of T with at least two linearly independent eigen vectors. If we write $f(X) = Xg(X)$ in $\mathbb{R}[X]$, then the linear operator $g(T)$ is zero.

Which of the following is true?

- (1) Both (a) and (b) are true.
- (2) Both (a) and (b) are false.
- (3) (a) is true and (b) is false.
- (4) (a) is false and (b) is true.

199. Let $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$ denote vectors in \mathbb{R}^n for a fixed $n \geq 2$. Which of the following defines an inner product on \mathbb{R}^n ?

- (1) $\langle x, y \rangle = \sum_{i,j=1}^n x_i y_j$
- (2) $\langle x, y \rangle = \sum_{i,j=1}^n (x_i^2 + y_j^2)$
- (3) $\langle x, y \rangle = \sum_{j=1}^n j^3 x_j y_j$
- (4) $\langle x, y \rangle = \sum_{j=1}^n x_j y_{n-j+1}$

200. Consider the quadratic form $Q(x, y, z)$ associated to the matrix

$$B = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & -2 \end{bmatrix}$$

Let

$$S = \left\{ \begin{bmatrix} a \\ b \\ c \end{bmatrix} \in \mathbb{R}^3 \mid Q(a, b, c) = 0 \right\}$$

Which of the following statements is FALSE?

- (1) The intersection of S with the xy-plane is a line
 (2) The intersection of S with the xz-plane is an ellipse
 (3) S is the union of two planes
 (4) Q is a degenerate quadratic form
- 201.** Let $l \geq 1$ be a positive integer. What is the dimension of the \mathbb{R} -vector space of all polynomials in two variables over \mathbb{R} having a total degree of at most l ?
- (1) $l+1$ (2) $l(l-1)$
 (3) $l(l+1)/2$ (4) $(l+1)(l+2)/2$
- 202.** Let A be a 3×3 matrix with real entries. Which of the following assertions is FALSE?
- (1) A must have a real eigenvalue
 (2) If the determinant of A is 0, then 0 is an eigenvalue of A
 (3) If the determinant of A is negative and 3 is an eigenvalue of A, then A must have three real eigenvalues
 (4) If the determinant of A is positive and 3 is an eigenvalue of A, then A must have three real eigenvalues
- 203.** Let A be a 3×3 real matrix whose characteristic polynomial $p(T)$ is divisible by T^2 . Which of the following statements is true?
- (1) The eigenspace of A for the eigenvalue 0 is two-dimensional
 (2) All the eigenvalues of A are real
 (3) $A^3 = 0$
 (4) A is diagonalizable

PART - C

- 204.** Let V be the vector space of all polynomials in one variable of degree at most 10 with real coefficients. Let W_1 be the subspace of V consisting of

polynomials of degree at most 5 and let W_2 be the subspace of V consisting of polynomials such that the sum of their coefficients is 0. Let W be the smallest subspace of V containing both W_1 and W_2 . Which of the following statements are true?

- (1) The dimension of W is at most 10
 (2) $W = V$
 (3) $W_1 \subset W_2$
 (4) The dimension of $W_1 \cap W_2$ is at most 5

- 205.** Let V be a finite dimensional real vector space and T_1, T_2 be two nilpotent operators on V. Let $W_1 = \{v \in V : T_1(v) = 0\}$ and $W_2 = \{v \in V : T_2(v) = 0\}$. Which of the following statements are FALSE?

- (1) If T_1 and T_2 are similar, then W_1 and W_2 are isomorphic vector spaces
 (2) If W_1 and W_2 are isomorphic vector spaces, then T_1 and T_2 have the same minimal polynomial
 (3) If $W_1 = W_2 = V$, then T_1 and T_2 are similar
 (4) If W_1 and W_2 are isomorphic, then T_1 and T_2 have the same characteristic polynomial

- 206.** Consider the following quadratic forms over \mathbb{R}

- (a) $6X^2 - 13XY + 6Y^2$,
 (b) $X^2 - XY + 2Y^2$,
 (c) $X^2 - XY - 2Y^2$.

Which of the following statements are true?

- (1) Quadratic forms (a) and (b) are equivalent
 (2) Quadratic forms (a) and (c) are equivalent
 (3) Quadratic form (b) is positive definite
 (4) Quadratic form (c) is positive definite

- 207.** Suppose A is a 5×5 block diagonal real matrix with diagonal blocks

$$\begin{pmatrix} e & & & & \\ & 1 & & & \\ & & e & & \\ & & & e & \\ & & & & e \end{pmatrix}, \begin{pmatrix} e & 1 & 0 \\ 0 & e & 0 \\ 0 & 0 & e \end{pmatrix}$$

Which of the following statements are true?

- (1) The algebraic multiplicity of e in A is 5
 (2) A is not diagonalizable
 (3) The geometric multiplicity of e in A is 3
 (4) The geometric multiplicity of e in A is 2

208. Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be a linear transformation satisfying $T^3 - 3T^2 = -2I$, where $I : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is the identity transformation. Which of the following statements are true?

- (1) \mathbb{R}^3 must admit a basis B_1 such that the matrix of T with respect to B_1 is symmetric.
- (2) \mathbb{R}^3 must admit a basis B_2 such that the matrix of T with respect to B_2 is upper triangular.
- (3) \mathbb{R}^3 must contain a non-zero vector v such that $Tv = v$.
- (4) \mathbb{R}^3 must contain two linearly independent vectors v_1, v_2 such that $Tv_1 = v_1$ and $Tv_2 = v_2$.

209. Let B be a 3×5 matrix with entries from \mathbb{Q} . Assume that $\{v \in \mathbb{R}^5 \mid Bv = 0\}$ is a three-dimensional real vector space. Which of the following statements are true?

- (1) $\{v \in \mathbb{Q}^5 \mid Bv = 0\}$ is a three-dimensional vector space over \mathbb{Q} .
- (2) The linear transformation $T : \mathbb{Q}^3 \rightarrow \mathbb{Q}^5$ given by $T(v) = B^t v$ is injective
- (3) The column span of B is two-dimensional
- (4) The linear transformation $T : \mathbb{Q}^3 \rightarrow \mathbb{Q}^3$ given by $T(v) = BB^t v$ is injective

210. Let V be the real vector space of real polynomials in one variable with degree less than or equal to 10 (including the zero polynomial). Let $T : V \rightarrow V$ be the linear map defined by $T(p) = p'$, where p' denotes the derivative of p . Which of the following statements are correct?

- (1) $\text{rank}(T) = 10$
- (2) $\text{determinant}(T) = 0$
- (3) $\text{trace}(T) = 0$
- (4) All the eigenvalues of T are equal to 0

211. Let V be an inner product space and let $v_1, v_2, v_3 \in V$ be an orthogonal set of vectors. Which of the following statements are true?

- (1) The vectors $v_1 + v_2 + 2v_3, v_2 + v_3, v_2 + 3v_3$ can be extended to a basis of V
- (2) The vectors $v_1 + v_2 + 2v_3, v_2 + v_3, v_2 + 3v_3$ can be extended to an orthogonal basis of V
- (3) The vectors $v_1 + v_2 + 2v_3, v_2 + v_3, 2v_1 + v_2 + 3v_3$ can be extended to a basis of V
- (4) The vectors $v_1 + v_2 + 2v_3, 2v_1 + v_2 + v_3, 2v_1 + v_2 + 3v_3$ can be extended to a basis of V

DECEMBER-2023**PART - B**

212. For $a \in \mathbb{R}$, let

$$A_a = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & a \end{pmatrix}. \text{ Which one of the}$$

following statements is true?

- (1) A_a is positive definite for all $a < 3$.
- (2) A_a is positive definite for all $a > 3$.
- (3) A_a is positive definite for all $a \geq -2$.
- (4) A_a is positive definite only for finitely many values of a .

213. We denote by I_n the $n \times n$ identity matrix. Which one of the following statements is true?

- (1) If A is a real 3×2 matrix and B is a real 2×3 matrix such that $BA = I_2$, then $AB = I_3$.

- (2) Let A be the real matrix $\begin{pmatrix} 3 & 3 \\ 1 & 2 \end{pmatrix}$. Then

there is a matrix B with integer entries such that $AB = I_2$.

- (3) Let A be the matrix $\begin{pmatrix} 3 & 1 \\ 1 & 2 \end{pmatrix}$ with

entries in $\mathbb{Z}/6\mathbb{Z}$. Then there is a matrix B with entries in $\mathbb{Z}/6\mathbb{Z}$ such that $AB = I_2$.

- (4) If A is a real non-zero 3×3 diagonal matrix, then there is a real matrix B such that $AB = I_3$.

214. Which one of the following statements is FALSE?

- (1) The product of two 2×2 real matrices of rank 2 is of rank 2.

- (2) The product of two 3×3 real matrices of rank 2 is of rank at most 2.
- (3) The product of two 3×3 real matrices of rank 2 is of rank at least 2.
- (4) The product of two 2×2 real matrices of rank 1 can be the zero matrix.

$$Q(x, y, z) = [x \ y \ z] A \begin{bmatrix} x \\ y \\ z \end{bmatrix} \text{ for all } x,$$

$y, z \in \mathbb{R}$ is invertible.

- 215.** Let $A = (a_{ij})$ be the $n \times n$ real matrix with $a_{ij} = ij$ for all $1 \leq i, j \leq n$. If $n \geq 3$, which one of the following is an eigenvalue of A ?

- (1) 1
- (2) n
- (3) $n(n+1)/2$
- (4) $n(n+1)(2n+1)/6$

- 216.** Let A be an $n \times n$ matrix with complex entries. If $n \geq 4$, which one of the following statements is true?

- (1) A does not have any non-zero invariant subspace in \mathbb{C}^n .
- (2) A has an invariant subspace in \mathbb{C}^n of dimension $n-3$.
- (3) All eigenvalues of A are real numbers.
- (4) A^2 does not have any invariant subspace in \mathbb{C}^n of dimension $n-1$.

- 217.** Let $(-, -)$ be a symmetric bilinear form on \mathbb{R}^2 such that there exists non-zero $v, w \in \mathbb{R}^2$ such that $(v, v) > 0 > (w, w)$ and $(v, w) = 0$. Let A be the 2×2 real symmetric matrix representing this bilinear form with respect to the standard basis. Which one of the following statements is true?

- (1) $A^2 = 0$.
- (2) $\text{rank } A = 1$.
- (3) $\text{rank } A = 0$.
- (4) There exists $u \in \mathbb{R}^2, u \neq 0$ such that $(u, u) = 0$.

PART - C

- 218.** Consider the quadratic form $Q(x, y, z) = x^2 + xy + y^2 + xz + yz + z^2$. Which of the following statements are true?

- (1) There exists a non-zero $u \in \mathbb{Q}^3$ such that $Q(u) = 0$.
- (2) There exists a non-zero $u \in \mathbb{R}^3$ such that $Q(u) = 0$.
- (3) There exist a non-zero $u \in \mathbb{C}^3$ such that $Q(u) = 0$.
- (4) The real symmetric 3×3 matrix A which satisfies

- 219.** Let \mathbb{F} be a finite field and V be a finite dimensional non-zero \mathbb{F} -vector space. Which of the following can NEVER be true?

- (1) V is the union of 2 proper subspaces.
- (2) V is the union of 3 proper subspaces.
- (3) V has a unique basis.
- (4) V has precisely two bases.

- 220.** Suppose a 7×7 block diagonal complex matrix A has blocks

$$(0), \quad (1), \quad \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \text{ and}$$

$$\begin{pmatrix} 2\pi i & 1 & 0 \\ 0 & 2\pi i & 0 \\ 0 & 0 & 2\pi i \end{pmatrix} \text{ along the diagonal.}$$

Which of the following statements are true?

- (1) The characteristic polynomial of A is $x^3(x-1)(x-2\pi i)^3$.
- (2) The minimal polynomial of A is $x^2(x-1)(x-2\pi i)^3$.
- (3) The dimensions of the eigenspaces for $0, 1, 2\pi i$ are 2, 1, 3 respectively.
- (4) The dimensions of the eigenspaces for $0, 1, 2\pi i$ are 2, 1, 2 respectively.

- 221.** Let $T : \mathbb{R}^5 \rightarrow \mathbb{R}^5$ be a \mathbb{R} -linear transformation. Suppose that $(1, -1, 2, 4, 0), (4, 6, 1, 6, 0)$ and $(5, 5, 3, 9, 0)$ span the null space of T . Which of the following statements are true?

- (1) The rank of T is equal to 2.
- (2) Suppose that for every vector $v \in \mathbb{R}^5$, there exists n such that $T^n v = 0$. Then T^2 must be zero.
- (3) Suppose that for every vector $v \in \mathbb{R}^5$, there exists n such that $T^n v = 0$. Then T^3 must be zero.
- (4) $(-2, -8, 3, 2, 0)$ is contained in the null space of T .

- 222.** Let X, Y be two $n \times n$ real matrices such that $XY = X^2 + X + I$. Which of the following statements are necessarily true?

- (1) X is invertible
- (2) X + I is invertible
- (3) XY = YX
- (4) Y is invertible

223. Consider $A = \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}$. Suppose $A^5 - 4A^4 - 7A^3 + 11A^2 - A - 10I = aA + bI$ for some $a, b \in \mathbb{Z}$. Which of the following statements are true?

- (1) $a + b > 8$
- (2) $a + b < 7$
- (3) $a + b$ is divisible by 2
- (4) $a > b$

224. Let A be an $n \times n$ real symmetric matrix. Which of the following statements are necessarily true?

- (1) A is diagonalizable.
- (2) If $A^k = I$ for some positive integer k, then $A^2 = I$.
- (3) If $A^k = 0$ for some positive integer k, then $A^2 = 0$.
- (4) All eigenvalues of A are real.

225. Let A be a real diagonal matrix with characteristic polynomial $\lambda^3 - 2\lambda^2 - \lambda + 2$. Define a bilinear form $\langle v, w \rangle = v^t A w$ on \mathbb{R}^3 . Which of the following statements are true?

- (1) A is positive definite.
- (2) A^2 is positive definite.
- (3) There exists a non-zero $v \in \mathbb{R}^3$ such that $\langle v, v \rangle = 0$.
- (4) rank A = 2.

LINEAR ALGEBRA
PREVIOUS YEAR PAPERS

ANSWERS

- | | | | | | |
|------------------|------------------|---------------|-------------------|----------------|----------------|
| 1. (2) | 2. (1) | 3. (2) | 157. (1, 2) | 158. (3) | 159. (3) |
| 4. (1) | 5. (3) | 6. (4) | 160. (2) | 161. (4) | 162. (4) |
| 7. (1, 3) | 8. (1, 2, 3) | 9. (2, 3, 4) | 163. (2) | 164. (2, 3) | 165. (1,2) |
| 10. (1, 2, 3, 4) | 11. (3, 4) | 12. (1, 2) | 166. (2, 4) | 167. (1, 3) | 168. (1, 2) |
| 13. (1, 3, 4) | 14. (2, 3) 1 | 5. (1, 3) | 169. (1, 2, 3, 4) | 170. (1,2,4) | 171. (2, 3) |
| 16. (2, 3) | 17. (1, 2) | 18. (1, 3, 4) | 172. (3) | 173. (3) | 174. (4) |
| 19. (1) | 20. (4) | 21. (2) | 175. (1) | 176. (2) | 177. (1) |
| 22. (1, 2) | 23. (2) | 24. (3) | 178. (1,2,3,4) | 179. (1,3) | 180. (2,4) |
| 25. (3) | 26. (1, 2) | 27. (1) | 181. (1,3) | 182. (4) | 183. (4) |
| 28. (1, 2) | 29. (1, 3, 4) | 30. (1, 2) | 184. (2,3,4) | 185. (2) | 186. (3) |
| 31. (2, 3) | 32. (1, 3, 4) | 33. (1, 2) | 187. (4) | 188. (4) | 189. (1) |
| 34. (1, 2) | 35. (3, 4) | 36. (3, 4) | 190. (4) | 191. (1) | 192. (3) |
| 37. (4) | 38. (2) | 39. (1) | 193. (4) | 194. (1,3,4) | 195. (1,2) |
| 40. (4) | 41. (4) | 42. (4) | 196. | 197. (2,4) | 198. (4) |
| 43. (1, 3) | 44. (2, 4) | 45. (1, 3) | 199. (3) | 200. (2) | 201. (4) |
| 46. (1, 3, 4) | 47. (1, 2, 3, 4) | 48. (1,2,3,4) | 202. (4) | 203. (2) | 204. (2, 4) |
| 49. (1) | 50. (2) | 51. (2) | 205. (2) | 206. (2,3) | 207. (1,2,3) |
| 52. (1) | 53. (3) | 54. (4) | 208. (1,2) | 209. (1,3) | 210. (1,2,3,4) |
| 55. (1, 2) | 56. (3, 4) | 57. (2, 3) | 211. (*) | 212. (2) | 213. (3) |
| 58. (2, 3, 4) | 59. (3) | 60. (2) | 214. (3) | 215. (4) | 216. (2) |
| 61. (1, 2, 3, 4) | 62. (1, 2) | 63. (1, 2) | 217. (4) | 218. (3,4) | 219. (1) |
| 64. (1) | 65. (1) | 66. (3) | 220. (1,4) | 221. (1,3,4) | 222. (1,3) |
| 67. (2) | 68. (4) | 69. (3) | 223. (2,3) | 224. (1,2,3,4) | 225. (2,3) |
| 70. (1) | 71. (1) | 72. (1, 4) | | | |
| 73. (1, 2, 3, 4) | 74. (3, 4) | 75. (1, 3) | | | |
| 76. (1) | 77. (2, 3) | 78. (2, 3) | | | |
| 79. (2) | 80. (1) | 81. (2) | | | |
| 82. (2) | 83. (4) | 84. (3) | | | |
| 85. (2, 4) | 86. (2, 3) | 87. (2, 4) | | | |
| 88. (1, 3, 4) | 89. (1, 2, 4) | 90. (1, 3) | | | |
| 91. (1, 4) | 92. (1, 4) | 93. (1) | | | |
| 94. (4) | 95. (4) | 96. (4) | | | |
| 97. (3) | 98. (1, 4) | 99. (3, 4) | | | |
| 100. (2, 3) | 101. (1, 4) | 102. (1,2,3) | | | |
| 103. (1, 2, 3) | 104. (3) | 105. (4) | | | |
| 106. (3) | 107. (4) | 108. (3) | | | |
| 109. (4) | 110. (2, 4) | 111. (4) | | | |
| 112. (3, 4) | 113. (1, 4) | 114. (4) | | | |
| 115. (1, 4) | 116. (2, 3) | 117. (3, 4) | | | |
| 118. (1) | 119. (3) | 120. (3) | | | |
| 121. (4) | 122. (2) | 123. (2) | | | |
| 124. (1, 4) | 125. (1, 3) | 126. (1, 4) | | | |
| 127. (3, 4) | 128. (4) | 129. (1, 4) | | | |
| 130. (4) | 131. (2) | 132. (2) | | | |
| 133. (3) | 134. (1) | 135. (4) | | | |
| 136. (3) | 137. (3) | 138. (2, 3) | | | |
| 139. (4,3) | 140. (1, 3) | 141. (1) | | | |
| 142. (4) | 143. (1, 2, 4) | 144. (2, 3) | | | |
| 145. (1, 3, 4) | 146. (4) | 147. (3) | | | |
| 148. (3) | 149. (4) | 150. (3) | | | |
| 151. (3) | 152. (2, 3) | 153. (3, 4) | | | |
| 154. (2, 3) | 155. (2, 4) | 156. (2) | | | |